

31 January 2017
EMA/HMPC/80629/2016
Committee on Herbal Medicinal Products (HMPC)

List of references supporting the assessment of *Salix* [various species including *S. purpurea* L., *S. daphnoides* Vill., *S. fragilis* L.], cortex

Final

The European Medicines Agency acknowledges that copies of the underlying works used to produce this monograph were provided for research only with exclusion of any commercial purpose.

Adamkiewicz VM, Fortier AA. Passage of salicin and saligenin across the wall of the rat ileum. *Can J Biochem Physiol* 1961, 39:1097-99

Akao T, Yoshino T, Kobashi K, Hattori M. Evaluation of salicin as an antipyretic prodrug that does not cause gastric injury. *Planta Med* 2002, 68:714-18

Albrecht M, Nahrstedt, A. Luepke, N.P. Theisen, N.L. Baron, G. Anti-inflammatory activity of flavonol glycosides and salicin derivatives from the leaves of *Populus tremuloides*. *Planta Med* 1990, 56:660

American Herbal Pharmacopoeia. Willow bark. Scotts Valley 1999

Aronson JK (Ed). Meyler's side effects of drugs. *Elsevier Science*, Amsterdam 2006

Baker S, Thomas PS. Herbal medicine precipitating massive haemolysis [letter]. *Lancet* 1987, 1:1039-40

Barnes J, Anderson LA, Phillipson JD. Herbal Medicines, 3rd edition. *Pharmaceutical Press*. London 2007, 598-600

Beer A-M, Loew D. Phytopharmaka und Rückenschmerzen. *Fortschritte der Medizin* 2008, 150:33-37

Beer A-M, Helleman M, Schübel H, Wegener T. Weidenrinde Extract bei Gon- und Coxarthrose. Zeitschrift für. *Phytotherapie* 2008, 29:215-222

Beer A-M, Wegener T. Willow bark extract (Salicis cortex) for gonarthrosis and coxarthrosis – Results of a cohort study with a control group. *Phytomedicine* 2008, 15:907-913

Benedum J, Loew D, Schilcher H. Arzneipflanzen in der traditionelle Medizin. Kooperation Phytopharmaka, Bonn. *Salicis cortex Monographie* 2006, 179:320-321



Biegert C, Wagner I, Lüdtke R, Kötter I, Lohmüller C, Günaydin I et al. Efficacy and safety of willow bark extract in the treatment of OA and RA: results of 2 randomised db controlled trials. *J Rheumatol* 2004, 31:2121-30

Bisset NG. *Herbal Drugs and Phytopharmaceuticals*. Medpharm Scientific Publishers, Stuttgart 1994, 437-439

Blaschek et al. In Hagers Handbuch der Pharmazeutischen Praxis. Folgenband 3. Drogen L-Z. Springer Verlag, Berlin Heidelberg 1998, 469-96

Bogduk N. Pharmacological alternatives for the alleviation of back pain. *Expert Opin Pharmacother* 2004, 5(10):2091-98

Boullata J, McDonnell PJ, Oliva CD. Anaphylactic reaction to a dietary supplement containing willow bark. *Ann Pharmacotherap* 2003, 37:832-35

BHMA. *British Herbal Pharmacopoeia*. Bournemouth 1983, 184-85

BHMA. *British Herbal Compendium*. Bournemouth 1992, 224-26

Bruneton J. Phytothérapie: les données de l'évaluation. *Tec et Doc Ed médicales internationales*. Paris 2002, 99-100

Cheng GF, Liu DP, Yang DX, He KQ, Bai JY, Zhu XY. Anti-inflammatory effects of tremulacin, a salicin-related substance isolated from *Populus tomentosa* Carr leaves. *Phytomedicine* 1994, 1:209-11

Chrubasik S, Eisenberg E, Balan E, Weinberger T, Luzzati R, Conradt C., Treatment of low back pain exacerbations with willow bark extract: a randomized double-blind study. *Am J Med* 2000, 109:9-14

Chrubasik S, Künzel O, Model A, Conradt C, Black A. Treatment of low back pain with a herbal or synthetic anti-rheumatic: a randomized controlled study. Willow bark extract for low back pain. *Rheumatology* 2001a, 40:1388-93

Chrubasik S, Kunzel O, Black A, Conradt C, Kerschbaumer F. Potential economic impact using a proprietary willow bark extract in outpatient treatment of low back pain: an open non-randomized study. *Phytomedicine* 2001b, 8(4):241-51

Chrubasik S, Pollak S. Wirksamkeit von Weidenrindeextrakt bei Schmerzen. *Z Phytother* 2002, 23:263-66

Chrubasik S. Weidenrindeextrakt verstärkt nicht die Regelblutung. *Naturamed* 2002, 17(12):13

Chrubasik S. Effects of willow bark and therapeutic activity. *Clin Pharmacol Ther*. 2003, 74:95-97

Chrubasik JE, Roufogalis BD, Chrubasik S. Evidence of effectiveness of herbal anti-inflammatory drugs in the treatment of painful osteoarthritis and chronic low back pain. *Phytother Res* 2007, 21:675-83

Clauson KA, Santamarina ML, Buettner CM, Cauffield JS. Evaluation of presence of aspirin-related warnings with willow bark. *Ann Pharmacother* 2005, 39:1234-37

Commission E. *Salicis cortex* (Weidenrinde). BAnz N° 228, 1984

CPMP/EMEA. Points to consider on clinical investigation of medicinal products used in the treatment of osteoarthritis. CPMP/EWP/784/97, 1998

CPMP/EMEA. Points to consider on clinical investigation of medicinal products other than NSAIDS for treatment of rheumatoid arthritis. CPMP/EWP/556/95 rev 1, 2003

DAB 10 Deutsche Apotheker Verlag, Stuttgart 1991

DAB 10. Kommentar. Deutsche Apotheker Verlag, Stuttgart 1995

Dingerman T, Hänsel R, Zündorf I. Pharmazeutische Biologie. Springer Verlag, Berlin 2002, 65-78

Drummond EM, Harbourne N, Marete E, Martyn D, Jacquier JC, O'Riordan D, Gibney ER. Inhibition of pro-inflammatory biomarkers in THP1 macrophages by polyphenols derived from chamomile, meadowsweet and willow bark. *Phytotherapy Research* 2013, 27: 588-596

Durak A, Gawlik-Dziki U. The Study of Interactions between Active Compounds of Coffee and Willow (*Salix* sp.) Bark Water Extract. *BioMed Research International* Volume 2014, Article ID 386953, 11 pages <http://dx.doi.org/10.1155/2014/386953>, 2014

EDQM Willow bark dry extract. *Pharmeuropa* 2006, 18 (1): 158-60

EDQM. European Pharmacopoeia 5th edition, Willow bark. *Pharmeuropa* 2006, 18 (3): 497-99

Enayat S, Banerjee S. The Ethanolic Extract of Bark from *Salix Aegyptiaca* L. Inhibits the Metastatic Potential and Epithelial to Mesenchymal Transition of Colon Cancer Cell Lines. *Nutrition and Cancer* 2014, 66(6): 999-1008

ESCOP. Monographs; second edition: *Salicis cortex*, 2003. 445-51

Esatbeyoglu T, Wray V, Winterhalter P. Dimeric Procyanidins: Screening for B1 to B8 and Semisynthetic Preparation of B3, B4, B6, and B8 from a Polymeric Procyanidin Fraction of White Willow Bark (*Salix alba*). *J. Agric. Food Chem.* 2010, 58: 7820-7830

Fiebig BL, Appel K. Anti-inflammatory effects of willow bark extract. *Clinical pharmacology & therapeutics* 2003, 96

Fiebig BL, Müller J, Kelber O, Okpanyi S, Weiser D, Appel K. Does willow bark have a sildenafil-like action? Effect of different fraction on the synthesis of c-GMP and NO in human umbilical vein endothelial cell (HUEC). *Planta Medica* 2010, 76: 1190

Fiebich BL, Chrubasik S. Effects of an ethanolic *Salix* extract on the release of selected inflammatory mediators *in vitro*. *Phytomed* 2004, 11: 135-38

Förster N, Ulrichs C, Zander M, Kätsel R, Mewis I. Salicylatreiche Weiden für die Arzneimittelherstellung. *Gesunde Pflanzen* 2009, 61: 129-134

Förster N, Ulrichs C, Zander M, Kätsel R, Mewis I. Factors influencing the variability of antioxidative phenolic glycosides in *Salix* species. *Agricultural and food chemistry* 2010; 58: 8205-8210. Fötsch G, Gründemann E, Pfeifer S, Hiller K, Salzwedel D. Biotransformation der Phenolglycosid Leiocarposid und Salicin. *Pharmazie* 1989a, 44: 555-58

Fötsch G, Pfeifer S. Die Biotransformation der Phenolglycosid Leiocarposid und Salicin – Beispiele für Besonderheiten von Absorption und Metabolismus glycosidischer Verbindungen. *Pharmazie* 1989b, 44: 710-2

Fötsch G, Pfeifer S. Vergleichende Serumspiegel-untersuchung von Salicylsäure nach oraler Applikation von Salicin bzw Natriumsalicylat in Ratten. *Pharmazie* 1990, 45: 535-36

Freischmidt A, Jürgenliemk G, Kraus B, Okpanyi SN, Müller J, Kelber O, et al. Contribution of flavonoids to the reduction of ICAM-1 expression in endothelial cells by a standardized willow bark extract. *Phytomedicine* 2012, 19: 245-252

Freischmidt A, Untergehrer M, Ziegler M, et al. Quantitative analysis of flavanones and chalcones from willow bark. *Pharmazie* 2015, 70: 565-568

Gagnier JJ, van Tulder MW, Berman B, Bombardier C. Herbal medicine for low back pain: a Cochrane Review. *Spine* 2007, 32(1):82-92

Gagnier JJ, Oltean H, MPH, _van Tulder MW, Berman BM, Bombardier C, Robbins CB. Herbal Medicine for Low Back Pain. *Spine* 2016, 41:116–133

Gawlik-Dziki U, Sugier D, Dziki D, Sugier P. Bioaccessibility *In Vitro* of Nutraceuticals from Bark of Selected *Salix* Species. Hindawi Publishing Corporation Scientific World Journal Volume 2014, Article ID 782763, 10 pages <http://dx.doi.org/10.1155/2014/782763>

Gopalan V, Pastuszyn A, Galey WR Jr, Glew RH. Exolytic hydrolysis of toxic plant glucosides by guinea pig liver cytosolic beta-glucosidase. *J Biol Chem* 1992, 267:14027-32

Hänsel. Weidenrinde, Salicis cortex. In: Phytopharmaka, 2nd ed. Berlin-Heidelberg-New York: Springer-Verlag 1991, 96-97

HagerRom. *Salicis cortex (Salix purpurea-Rinde)*, 2001

Hostanska K, Jürgenliemk G, Abel G, Nahrstedt A, Saller R. Willow bark extract (BNO 1455) and its fractions suppress growth and induce apoptosis in human colon and lung cancer cells. *Cancer Detection and Prevention* 2007, 31:129-131

Ishikado A, Sono Y, Matsumoto M, et al. Willow bark extract increases antioxidant enzymes and reduces oxidative stress through activation of Nrf2 in vascular endothelial cells and *Caenorhabditis elegans*. *Free Radical Biology and Medicine* 2013, 65:1506–1515

Julkunen-Tiitto R, Meier B. The enzymatic decomposition of salicin and its derivatives obtained from *Salicaceae* species. *J Nat Prod* 1992a, 55:1204-12

Julkunen-Tiutto R, Gebhardt K. Further studies on the drying of willow twigs: the effect of low drying temperature on labile phenolics. *Planta Med* 1992b, 58: 385-86

Julkunen-Tiutto R, Sorsa S. Testing the effects of drying methods on willow flavonoids, tannins and salicylates. *J Chem Ecol* 2001, 27(4):779-89

Kammerer B, Kahlich R, Biegert C, Gleiter CH, Heide L. HPLC-MS/MS analysis of willow bark extracts contained in pharmaceutical preparations. *Phytochem Anal* 2005, 16:470-78

Kaufeld AM, Pertz HH, Kolodziej H. A Chemically Defined 2,3-trans Procyanidin Fraction from Willow Bark Causes Redox-Sensitive Endothelium-Dependent Relaxation in Porcine Coronary Arteries. *J. Nat. Prod.* 2014, 77:1607–1614

Kaul R, Lagoni N. Weidenrinde. Renaissance eines Phytoanalgetikums. *Deutsche Apothekerzeitung* 1999, 19(37):43-50

Kelber O, Bonnaterre G, Kinscherf R, Weiser D, Metz J. Inhibitorische Effekte von Weidenrindenextrakten auf proinflammatorische Prozesse in LPS aktivierten Humanmonozyten. *Zeitschrift für Rheumatologie* 2006, 65 Suppl 1:S31

Kelber O, Okpanyi SN, Freischmidt A, et al. Antidepressant mechanism of action of willow bark extract STW I-33. *Planta Medica* 2011, 77:1444

Keusgen M, Algauer-Lechner C. Vielstoffgemisch zwischen Entzündungen und Schmerzen. *Pharmazeutische Zeitung* 2007, 152:16-22

Khayyal MT, El-Ghazaly MA, Abdallah DM, Okpanyi SN, Kelber O, Weiser D. Mechanisms involved in the anti-inflammatory effect of a standardised willow bark extract. *Arzneim. Forsch / Drug Research* 2005, 55(11):677-87

- Kim CS, Subedi L, Jin Park K, et al. Salicin derivatives from *Salix glandulosa* and their biological activities. *Fitoterapia* 2015, 106:147–152
- Knuth S, Abdelsalam RM, Khayyal MT, et al. Catechol conjugates are *in vivo* metabolites of *Salicis cortex*. *Planta Med* 2013, 79:1489–1494
- Kong CS, Kim KH, Choi JS, Kim JE, Park C, Jeong JW. Salicin, an Extract from White Willow Bark, inhibits Angiogenesis by Blocking the ROS-ERK Pathways. *Phytother. Res.* 2014, 28:1246–1251
- Krivoy N, Pavlotzky E, Chrubasik S, Eisenberg E, Brook G. Effect of *Salicis cortex* extract on human platelet aggregation. *Planta Med* 2001, 67:209-12
- Kuppsamy UR, Khoo HE, Das NP. Structure-activity studies of flavonoids as inhibitors of hyaluronidase. *Biochem Biophys* 1990, 140:397-401
- Lardos A, Schmidlin C.B., Fischer M., Ferlas-Chlodny E., Loniewski I., Samochowiec L., et al. Wirksamkeit und Verträglichkeit eines wässrig ausgezogenen Weidenrindenextraktes bei Patienten mit Hüft-und Kniearthrose. *Z Phytotherap* 2004, 25:275-81
- Leporatti ML, Pavesi A. New or uncommon uses of several medicinal plants in some areas of central Italy. *J Ethnopharmacol* 1990, 29:213-23
- Madaus A. Lehrbuch biologische Heilmittel 1935, 2394-2399
- März RW, Kemper F. Weidenrindenextrakt – Wirkungen und Wirksamkeit. Erkenntnisstand zur Pharmacologie, Toxikologie und Klinik. Wien. Med. *Wochenschrift* 2002, 152:354-359
- Matsumoto Y, Ohsako M, Takadate A, Goto S. Reduction of erythrocyte membrane permeability and protein binding of LMW drugs following glycoside derivatisation. *J Pharm Sci* 1993, 82:399-403.
- Mayer RA, Mayer M. Biologische Salicyltherapie mit Cortex Salicis (Weidenrinde). *Pharmazie* 1949, 4:77-81
- McGuffin M, Hobbs C, Upton R. Botanical Safety Handbook. CRC Press Boca Raton Florida 1997 , 154-155
- Meier B, Lehmann D, Sticher O, Bettschart A . Identification and determination of 8 phenol glycosides each in *Salix purpurea* and *S. daphnoides* by modern HPLC. *Pharm Act Helv* 1985a, 60:269-75
- Meier B, Sticher O, Zürich, Bettschart A Weidenrinde – Qualität. Gesamtsalicinbestimmung in Weidenrinden und Weidenpräparaten mit HPLC. *Deutsche Apoth Ztg* 1985b, 125:341-47
- Meier B, Libi M. Salizinhaltige pflanzliche Arzneimittel. Überlegungen zur Wirksamkeit und Unbedenklichkeit. *Z Phytother* 1990, 11:50-58
- Mills SY, Jacoby RK, Chacksfield M, Willoughby M. Effect of a proprietary herbal medicine on the relief of chronic arthritic pain: a double-blind study. *Br J Rheumatol* 1996, 35:874-78
- Morgan M, Bone K, Mills S, McMillan J. Part 2: Safety Monographs, in: Mills S and Bone K, The Essential Guide Herbal Safety, Elsevier Churchill Livingstone, St. Louis 2005, 626-628
- Moro PA, Flacco V, Cassetti F, Clementi V, Colombo ML, Chiesa GM, Menniti-Ippolito F, Raschetti R, Santucci C. Hypovolemic shock due to severe gastrointestinal bleeding in a child taking an herbal syrup. *Ann Ist Super Sanità*, 2011, Vol. 47, 3:278-283
- Müller-Fassbender H, Müller J, Bach G. Wirksamkeit und Verträglichkeit eines Weidenrindenpräparates bei Erkrankungen des Bewegungsapparates. *Arthritis und Rheuma* 2007, 27:267-271

Müller J, Kelber O, Weiser D, Stange S. Willow bark extract STW 33-1 in the long term treatment of osteoarthritis and low back pain. *Planta Medica* 2010, 76:1361

Oltean H, Robbins C, van Tulder MW, Berman BM, Bombardier C, Gagnier JJ. Herbal medicine for low back pain. Cochrane Database of Systematic Reviews 2014, Issue 12. Art. No.: CD004504 DOI: 10.1002/14651858.CD004504.pub4

Nahrstedt A, Schmidt M, Jäggi R, Metz J, Khayyal MT. Willow bark extract: the contribution of polyphenols to the overall effect. *Wien Med Wochenschr* 2007, 157(13-14):348-51

Pentz R, Busse HG, König R, Siegers CP. Bioverfügbarkeit von Salicylsäure und Coffein aus einem phytoanalgetischen Kombinationspräparat. *Z Phytother* 1989, 10: 92-96

Pugliese A, Beltramo T, Torre D. Reye's and Reye's-like syndromes. *Cell Biochem Function* 2008; 26: 741-746

Rice-Evans CA, Miller NJ, Bolwell PG, Bramley PM, Pridham JB. The relative antioxidant activities of plant-derived polyphenolic flavonoids. *Free Rad Ser* 1995, 22(4):375-83

Rohnert U, Koske D, Schneider W, Elstner EF. Inhibition by *Salix*-extracts and Phytodolor® of copper-catalyzed oxidative destructions. *Z Naturforsch* 1998, 53:233-40

Rotblatt M. Evidence-based herbal medicine. Hanley and Belfus, Philadelphia 2002, 360-63

Saller R, Melzer J, Felder M, the Swiss Assalix Study Group. Pain relief with a proprietary extract of willow bark in rheumatology. An open trial. *Schweiz Zschr GanzheitsMedizin* 2008, 20(3):156-62

Samochowiec L. Efficacy of Cortex *Salix* Extract (*Salix* (r)) in patients with arthrosis in double blind, randomized, verum controlled clinical study 2001 (unpublished)

Schmid B, Kötter I, Heide L. Pharmacokinetics of salicin after oral administration of a standardized willow bark extract. *Eur J Clin Pharmacol* 2001a, 57:387-91

Schmid B, Lüdtke R, Selbmann HK, Kötter I, Tschirdehn B, Schaffner W, et al. Efficacy and tolerability of a standardized willow bark extract in patients with osteoarthritis: randomized placebo-controlled double blind clinical trial. *Phyt Res* 2001b, 15: 344-50

Schmid B, Biegert C, Kötter I, Heide L. Pharmakinetik von Salicin nach oraler Gabe eines standardisierten Weidenrindenextraktes, in: *Phytopharmaka VII*. Schulz V, Rietbrock N, Roots I, Loew D, *Steinkopf*, Darmstadt 2002: 93-100

Setty AR, Sigal LH. Herbal medications commonly used in the practice of rheumatology: mechanisms of action, efficacy and side effects. *Semin Arthritis Rheum* 2005, 34: 773-84

Shalansky S, Lynd L, Richardson K, Ingaszewski A, Kerr C. Risk of warfarin-related bleeding events and supratherapeutic INRs associated with CAM: a longitudinal analysis. *Pharmacother* 2007, 27(9):1237-47

Shakibaei M, Allaway D, Nebrich S, Mobasher A. Botanical Extracts from Rosehip (*Rosa canina*), Willow Bark (*Salix alba*), and Nettle Leaf (*Urtica dioica*) Suppress IL-1 β -Induced NF- κ B Activation in Canine Articular Chondrocytes. *Evidence-Based Complementary and Alternative Medicine* Volume 2012, Article ID 509383, 16 pages doi:10.1155/2012/509383

Shara M, Stohs SJ. Efficacy and Safety of White Willow Bark (*Salix alba*) Extracts. *Phytother. Res.* 2015, 29:1112–1116

Shrivastava R, Pechadre JC, John GW. *Tanacetum parthenium* and *Salix alba* (Mig-RL) combination in migraine prophylaxis. *Clin Drug Invest* 2006, 26(5):287-96

SPC Aspirin Junior; May 2016

Stange R, Müller J, Kelber O, Uehleke B. Willow bark extract STW 33-I is safe and effective in the long term treatment of outpatients with chronic musculoskeletal pain. *J Alternative Complementary Med* 2014, 20:A83

Steinegger E, Hövel H. Analytische und biologische Untersuchungen an Salicaceen-Wirkstoffen, insbesondere Salicin. *Pharm Acta Helv* 1972, 47:222-34

Tunón H1, Olavsdotter C, Bohlin L. Evaluation of anti-inflammatory activity of some Swedish medicinal plants. Inhibition of prostaglandin biosynthesis and PAF-induced exocytosis. *J Ethnopharmacol* 1995, 48:61-76

Uehleke B, Preis S, Müller J, Stange R, Kelber O, Melzer J. Willow bark extract STW 33-I in the long-term treatment of outpatients with rheumatic pain mainly osteoarthritis or back pain. *Phytomedicine* 20 2013a, 980–984

Uehleke B, Müller J, Stange R, Kelber O, Melzer J. Willow bark extract STW 33-I is safe and effective in the long term treatment of outpatients with rheumatic pain esp. osteoarthritis or backpain - a subgroup analysis. *Planta Medica* 2013b, 79:1273 (abstract)

Ulrich-Merzenich G, Kelber O, Freischmidt A, et al. Evidences for an antidepressant effect of willow bark in the Porsolt swimming test. *European Journal of Integrative Medicine* 2010, 2:188 (abstract)

Ulrich-Merzenich G, Koptina A, Kelber O, et al. Comparative analysis of gene expression profiles from rats treated with a standardized willow bark preparation, its salicin rich ethanol fraction and imipramine for toxicological outcomes. *Planta Medica* 2011, 77:1251

Ulrich-Merzenich G, Koptina A, Kelber O, et al. Prediction of adverse events by in vivo gene expression profiling exemplified for phytopharmaceuticals containing salicylates and the antidepressant imipramine. *Phytomedicine* 2012, 19:322–329

Ulrich-Merzenich G, Hartbrod F, Koptina A. Synergie-Effekte als Grundprinzip der Wirkung von Phytopharmaka am Beispiel der Weidenrinde (*Salix* spec.) *Z Phytother* 2013, 34:34

Vlachojannis JE, Cameron M, Chubasick S. A systematic review on the effectiveness of willow bark for musculoskeletal pain. *Phytotherapy Research* 2009, 23:897-900

Vlachojannis J, Magora F, Chubasik S. Willow Species and Aspirin: Different Mechanism of Actions. *Phytother. Res.* 2011, 25:1102–1104

Volmink J. The willow as a Hottentot (Khoikhoi) remedy for rheumatic fever. *J R Soc Med* 2008, 101:321-323

Wagner I, Greim I, Laufer C, Heide S, L, Gleiter CH. Influence of willow bark extract on cyclooxygenase activity and on TNF alpha or interleukin 1 beta release in vitro and in vivo. *Clin Pharmacol Therap* 2003a, 73(3):272-74

Wagner I, Biegert C, und Lutz Heide, Aktuelle Forschungsergebnisse zur Weidenrinde. *Pharm. Ztg* 2003b, 138:21-32

Wagner I., Effects of willow bark. *Clin Pharmacol Therap* (letters to editor)2003c, 96-97

Wegener T. Zur Studienlage der Weidenrindenextrakten. *Zietschrift für Phytotherapie* 2009, 30:232-237

Werner G, Scheithe K. Willow bark extract (Assalix ®) for chronic back pain an arthralgia, a post-authorisation study. *Phytopharmaka und Phytotherapie* 2004, Forschung und Praxis, Berlin (poster abstract)

Wichtl M (ed) Teedrogen und Phytopharmaka 4th ed, Stuttgart: *Wissenschaftliche Verlagsgesellschaft*, 2002, 534-7

Wiesneth S, Petereit F, Jürgenliemk G. *Salix daphnoides*: a screening for oligomeric and polymeric proanthocyanidins. *Molecules* 2015, 20:13764-13779

Williamson E, Driver S, Baxter K. Stockley's herbal medicines interactions. *Pharmaceutical Press* 2nd ed. London 2013, 485

Wuthold K, Germann I, Roos G, Kelber O, Weiser D, Heinle H, Kovar KA. TLC and multivariate data analysis of willow bark extracts. *J Chromatogr Sci* 2004, 42(6):306-09

Wutzke KD. Radioisotopenmarkierung und Pharmakokinetik von Phenolglycosiden. Dissertation Phillipsuniversität Marburg 1991