

21 November 2017
EMA/HMPC/325719/2017
Committee on Herbal Medicinal Products (HMPC)

List of references supporting the assessment of *Hedera helix L., folium*

Final

The European Medicines Agency acknowledges that copies of the underlying works used to produce this monograph were provided for research only with exclusion of any commercial purpose.

AIFA (L'Agenzia Italiana Del Farmaco). I'Ufficio di Farmacovigilanza dell'Agenzia Italiana del Farmaco (AIFA). Nota informativa importante dell'agenzia italiana del farmaco (aifa). mucolitici per uso orale e rettale: controindicazione nei bambini al di sotto dei 2 anni. Novembre 2010a. Available at:
http://www.aifa.gov.it/sites/default/files/nii_mucolitici_aifa_versione_definitiva_17.11.2010_.pdf, Accessed: 17/09/2016

AIFA (L'Agenzia Italiana Del Farmaco). Domande e risposte. Informazioni sulla tosse nei bambini sotto i 2 anni d'età: perché i mucolitici non devono più essere utilizzati? 30 novembre 2010b. Available at:
http://www.aifa.gov.it/sites/default/files/domande_e_rispostemucoliticifinale22112010.pdf, Accessed: 17/09/2016

Amara-Mokrane YA, Lehucher-Michel MP, Balansard G, Dumenil G, Botta A. Protective effects of α-hederin, chlorophyllin, and ascorbic acid towards the induction of micronuclei by doxorubicin in cultured human lymphocytes. *Mutagenesis* 1996, 11(2):161-167

American Academy of Paediatrics. Committee on Drugs. Use of Codeine- and Dextromethorphan-Containing Cough Remedies in Children (RE9722). Policy statement. *Pediatrics* 1997, 99(6):5 pages. Available at: <http://host.acep.org/tmp3.secure-xp.net/aapacep/polstat/ucdccrc.pdf>

ANSM (L'Agence nationale de sécurité du médicament et des produits de santé). Commission Nationale de Pharmacovigilance. Enquête Officielle de Pharmacovigilance sur les effets indésirables des spécialités mucolytiques chez le nourrisson de moins de deux ans. In: Compte rendu de la réunion du mardi 24 mars 2009. Saint-Denis, le 19 mai 2009, p 17-20. Available at:
https://ansm.sante.fr/var/ansm_site/storage/original/application/7705f2db6f54e980bc0695c6ca8358a3.pdf, Accessed: 25/03/2017

ANSM (L'Agence nationale de sécurité du médicament et des produits de santé). Commission Nationale de Pharmacovigilance. Enquête Officielle de Pharmacovigilance sur les effets indésirables des



specialités mucolytiques chez le nourrisson de moins de deux ans. In: Compte rendu de la réunion du mardi 30 mars 2010. Saint-Denis, le 25 mai 2010a, p 13-21. Available at:
https://ansm.sante.fr/var/ansm_site/storage/original/application/7705f2db6f54e980bc0695c6ca8358a3.pdf, Accessed: 25/03/2017

ANSM (L'Agence nationale de sécurité du médicament et des produits de santé). Nouvelles modalités de prise en charge de la toux chez le nourrisson (enfant de moins de 2 ans) - Point d'information. 28/10/2010b. Available at: <https://ansm.sante.fr/S-informer/Points-d-information-Points-d-information/Nouvelles-modalites-de-prise-en-charge-de-la-toux-chez-le-nourrisson-enfant-de-moins-de-2-ans-Point-d-information>, Accessed: 25/03/2017

Arch F. Erfahrungsberichte über die Aerosol-Behandlung der Bronchitis mit Prospan®. *Notabene Medici* 1974, 4(6):1-7

Balansard G, Timon-David P, Julien J, Bernard P, Gasquet M. Douvicultural and antifungal activities of α-Hederin extracted from *Hedera helix* L. leaves. *Planta Med* 1980, 39:234

Becker C. Wirkmechanismus von Efeu entschlüsselt. *Pharmazeutische Zeitung* 2003, 148(49):28

Beden AB, Perko J, Tercelj R, Kreft S. Treatment of acute respiratory infections in Slovenian children with ivy leaf extract syrup. *Zdrav Vestn* 2011, 80:276-84 [Slovenian; English abstract]

Beyer D. Zellbiologische Wirkmechanismen von Efeublattextrakt entschlüsselt. *Ärztezeitsch Naturheilverf* 2005, 46(3):167

Blaschek W, Ebel S, Hackenthal E, Holzgrabe U, Keller K, Reichling J, Schulz V, editors. HagerROM 2006: Hagers Handbuch der Drogen und Arzneistoffe. Version 5.0. Springer-Verlag, Heidelberg 2006

Blaschek W, Hilgenfeldt U, Holzgrabe U, Reichling J, Ruth P, Schulz V, editors. HagerROM 2014: Hagers Enzyklopädie der Arzneistoffe und Drogen. Version 10.6. Springer-Verlag, Heidelberg 2014

Böhlau V. Therapeutische Erfahrungen mit Prospan bei chronisch-obstruktiven Atemwegs-erkrankungen. *Notabene Med* 1977, 7(11):26-29

Bolbot Y, Prokhorov E, Mokha S, Yurtseva A. Comparing the efficacy and safety of high-concentrate (5-7.5:1) ivy leaves extract and Acetylcysteine for treatment of children with acute bronchitis. *Drugs Ukr* 2004, 11:1-4

Boll PM, Hansen L. On the presence of falcarinol in Araliaceae. *Phytochemistry* 1987, 26(11):2955-2956

Boyle J, Harman RMH. Contact dermatitis to *Hedera helix* (Common Ivy). *Contact Derm* 1985, 12:111-112

Brattström A, Schroder E, Oberst M. Wirksamkeit von Efeu Hustensirup Madaus und Prospan Hustensaft bei Kindern mit Bronchitis. *Kinder Jugendmed* 2006, 6:191-4

Bregnbak D, Menné T, Johansen JD. Airborne contact dermatitis caused by common ivy (*Hedera helix* L. ssp. *helix*). *Contact Derm* 2015, 72 (4):243-244

Brömel J, Zettl K. Efeuvergiftung bei Rehwild. *Der Praktische Tierarzt* 1986, 67:967-968

Bruhn G, Faasch H, Hahn H, Hausen BM, Bröhan J, König WA. Das Auftreten von Falcarinol und Didehydrofalcarinol in Efeu (*Hedera helix* L.). *Zeitschr Naturforsch* 1987, 42B:1328-1332

Brussell DE. Medicinal plants of Mt. Pelion, Greece. *Econ Bot* 2004, 58:174-202

Büechi S. Efeu. *Schweizer Apothekerzeitung* 2002, 140(20):700-701

Büechi S, Kähler D. Efeu (*Hedera helix*) bei Atemwegserkrankungen. *Schweizer Zeitschr Ganzheitsmed* 2003, 15(3):124-128

Bun SS, Elias R, Baghdikian B, Ciccolini J, Ollivier E, Balansard G. α -Hederin potentiates 5-FU antitumor activity in human colon adenocarcinoma cells. *Phytoth Res* 2008, 22:1299-1302

Cahiers de L'Agence N°3 - Médicaments à base de plantes - Ministry of Health and Family, France 1998; 45, 57, 59, 73 (note: former version: Bulletin official Nr. 90/22 1990)

Capasso A, Pinto A, Mascolo N, Autore G, Capasso F. Reduction of agonist-induced contractions of guinea-pig isolated ileum by flavonoids. *Phytother Res* 1991, 5(2):85-87

Centre for Reviews and Dissemination. Efficacy of dry extract of ivy leaves in children with bronchial asthma: a review of randomized controlled trials (structured abstract). *Database Abstracts of Reviews of Effects* 2008, 1-4

Chichiricco G, Cifani MP, Frizzi G, Tammaro F. Phytotherapy in the Subequana Valley, Abruzzo, central Italy. *J Ethnopharmacol* 1980, 2:247-257

CHMP. Codeine not to be used in children below 12 years for cough and cold. EMA/249413/2015, London 2015

Cioaca C, Margineanu C, Cucu V. The saponins of *Hedera helix* with antibacterial activity. *Pharmazie* 1978, 33(9):609-610

COPD-Clinical Practice Guideline. Global Strategy for the Diagnosis, Management, and Prevention of COPD. MedStar Health 2016, 1-8

Cwientzek U, Ottlinger B, Arenberger P. Acute bronchitis therapy with ivy leaves extracts in a two-arm study. A double-blind, randomised study vs. another ivy leaves extract. *Phytomedicine* 2011, 18:1105-1109

Czygan FC. *Hedera helix* L. - Der Efeu. *Zeitschr Phytother* 1990, 11:133-138

Danloy S, Quetin-Leclercq J, Coucke P, DePauw-Gillet MC, Elias R, Balansard G et al. Effects of α -hederin, a saponin extracted from *Hedera helix*, on cells cultured *in vitro*. *Planta Med* 1994, 60:45-49

Daston GP, Overmann GJ, Baines D, Taubeneck MW, Lehman-McKeeman LD, Rogers JM et al. Altered Zn status by α -hederin in the pregnant rat and its relationship to adverse developmental outcome. *Reprod Toxicol* 1994, 8(1):15-24

DEGAM-Leitlinie Nr. 11. Husten. Omicron publishing Düsseldorf 2008: 1-64 (http://www.degam.de/typo/uploads/media/LL-11_)

Delmas F, Di Giorgio C, Elias R, Gasquet M, Azas N et al. Antileishmanial activity of three saponins isolated from ivy, α -hederin, β -hederin and hederacolchiside A₁, as compared to their action on mammalian cells cultured *in vitro*. *Planta Med* 2000, 66:343-347

De Medeiros JMR, Macedo M, Contancia JP, Nguyen Chi, Cunningham G et al. Antithrombin activity of medicinal plants of the Azores. *J Ethnopharmacol* 2000, 72(1-2):157-165

De Smet PAGM. *Hedera helix*. In: de Smet P, Keller K, Hänsel R, editors. Adverse Effects of Herbal Drugs. Springer-Verlag Berlin, Heidelberg, New York 1993, 209-215

Dorsch W, Loew D, Meyer-Buchtela E, Schilcher H. *Hederae helicis folium* (Efeublätter) Kinderdosierungen von Phytopharmaaka, 3rd ed. Vol I. Kooperation Phytopharmaaka 2002, 72-73

Düchtel-Brühl Ä. Ergebnisse der Behandlung spastischer Bronchitiden im Kindesalter mit Prospan. *Med Welt* 1976, 27(10):481

Duffy JY, Baines D, Overmann GJ, Keen CL, Daston GP. Repeated administration of α -hederin results in alterations in maternal zinc status and adverse developmental outcome in the rat. *Teratology* 1997, 56: 327-334

Elias R, De Méo M, Vidal-Ollivier E, Laget M, Balansard G et al. Antimutagenic activity of some saponins isolated from *Calendula officinalis* L, *C. arvensis* L. and *Hedera helix* L. *Mutagenesis* 1990, 5(4):327-331

El-Marzabani MM, El-Aaser AA, Attia MA, El-Duweini AK, Ghazal AM. Screening system for Egyptian plants with potential anti-tumour activity. *Planta Med* 1979, 36:150-155

EMA. Points to consider on clinical investigation of medicinal products in the chronic treatment of patients with chronic obstructive pulmonary disease (COPD). EMEA (CPMP/EWP/562/98), London 1999

ESCOP Monographs 2nd ed. *Hederae helicis folium* – ivy leaf. European Scientific Cooperative on Phytotherapy, editor. Thieme, Stuttgart 2003, 241-247

European Pharmacopoeia 9.0th ed. Ivy leaf – *Hederae folium*. Council of Europe. 01/2017:2148

Facino RM, Carini M, Bonadeo P. Efficacy of topically applied *Hedera helix* L. saponins for treatment of liposclerosis (so-called "cellulitis"). *Acta Therap* 1990, 16: 337-349

Facino RM, Carini M, Stefani R, Aldini G, Saibene L. Anti-elastase and anti-hyaluronidase activities of saponins and sapogenins from *Hedera helix*, *Aesculus hippocastanum*, and *Ruscus aculeatus*: factors contributing to their efficacy in the treatment of venous insufficiency. *Archiv Pharm* (Weinheim) 1995, 328: 720-724

Favel A, Steinmetz MD, Ollivier E, Elias R, Régli P et al. Screening of triterpenoid saponins for antifungal activity. *Planta Med* 1992, 58(1):A635-636

Favel A, Steinmetz MD, Regli P, Vidal-Ollivier E, Elias R et al. In vitro antifungal activity of triterpenoid saponins. *Planta Med* 1994, 60:50-53

Fazio S, Pouso J, Dolinsky D, Fernandez A, Hernandez M, Clavier G et al. Tolerance, safety and efficacy of *Hedera helix* extract in inflammatory bronchial diseases under clinical practise conditions: a prospective, open, multicentre post-marketing study in 9657 patients. *Phytomedicine* 2009, 16(1):17-24

Frohne D, Pfänder HJ. *Hedera helix* L. Giftpflanzen, 5th ed. Wissenschaftliche Verlagsgesellschaft, Stuttgart 2004, 75-77

Fuchs J, Rauber-Luthy C, Kupferschmidt H, Kupper J, Kullak-Ublick G-A, Ceschi A. Acute plant poisoning: Analysis of clinical features and circumstances of exposure. *Clin Toxicol* 2011, 49(7):671-680

Gafner F, Epstein W, Reynolds G, Rodriguez E. Human maximization test of falcarinol, the principal contact allergen of English ivy and Algerian ivy (*Hedera helix*, *H. canariensis*). *Contact Derm* 1988, 19:125-128

Gaillard Y, Blaise P, Darré A, Barbier T, Pépin G. An unusual case of death: suffocation caused by leaves of common ivy (*Hedera helix*). Detection of hederacoside C, α -hederin, and hederagenin by LC-EI/MS-MS. *J Anal Toxicol* 2003, 27:257-262

García M, Fernández E, Navarro JA, del Pozo MD, Fernández de Corrès L. Allergic contact dermatitis from *Hedera helix* L. *Contact Derm* 1995, 33:133-134

Gepdiremen A, Mshvildadze V, Süleyman H, Elias R. Acute anti-inflammatory activity of four saponins isolated from ivy: α -hederin, hederasaponin-C, hederacolchiside-E and hederacolchiside-F in carrageenan-induced rat paw oedema. *Phytomedicine* 2005, 12:440-444

Goetz P. Plantes anti-adhésives d'helicobacter. *Phytotherapie* 2007, 5:101

Greunke C, Hage-Hülsmann A, Sorkalla T, Keksel N and Häberlein F. A systematic study on the influence of the main ingredients of an ivy leaves dry extract on the β_2 -adrenergic responsiveness of human airway smooth muscle cells. *Pulm Pharm & Therap* 2015, 31:92-98

Gülcin I, Mshvildadze V, Gepdiremen A, Elias R. Antioxidant activity of saponins isolated from ivy: α -hederin, hederasaponin-C, hederacolchiside-E and hederacolchiside-F. *Planta Med* 2004, 70(6):561-563

Gulyas A, Lämmlein MM. Zur Behandlung von Kindern mit chronisch-obstruktiver Bronchitis. *Sozialpädiatrie* 1992, 14(8):632-634

Gulyas A, Repges R, Dethlefsen U. Konsequente Therapie chronisch-obstruktiver Atemwegserkrankungen bei Kindern. *Atemwegs- Lungenkrank* 1997, 23(5):291-294

Gulyas A. Efeublätter-Extrakt zeigt bei Kindern mit chronisch obstruktiven Atemwegserkrankungen vergleichbare Effekte wie Acetylcystein. 1999, Available at: www.phytotherapie-komitee.de

Guo R, Pittler MH, Ernst E. Herbal medicines for the treatment of COPD: a systematic review. *Eur Respir J* 2006, 28(2):330-338

Haen E. Pharmacological activities of *Thymus vulgaris* and *Hedera helix*. *Phytomedicine* 1996, 3(1):144

Hänsel R, Keller K, Rimpler H, Schneider G, editors. *Hagers Handbuch der Pharmazeutischen Praxis*. 5th ed. Springer-Verlag, Berlin 1993, 398-407

Hänsel R, Sticher O. *Pharmakognosie-Phytopharmazie*. 7th ed. Springer-Verlag, Heidelberg 2004, 493

Hannu T, Kauppi P, Tuppurainen M, Piirilä P. Occupational asthma to ivy (*Hedera helix*). *Allergy* 2008, 63:482-3

Hausen BM, Bröhan J, König WA, Faasch H, Hahn H, Bruhn G. Allergic and irritant contact dermatitis from falcarinol and didehydrofalcarinol in common ivy (*Hedera helix* L.). *Contact Derm* 1987, 17:1-9

Hausen BM. Allergiepflanzen-Pflanzenallergene, Teil 1: Kontaktallergene, Ecomed Verlags-gesellschaft, Landsberg/Lech 1988, 147-149

Hecker M. Hustentropfen mit Ethanol - deutlich bessere Wirksamkeit. *T&E Pädiatrie* 1997a, 10:648-649

Hecker M. Verschiedene Zubereitungen von Efeublättertrockenextrakt: Dosisanpassung erforderlich. *T&E Pädiatrie* 1997b, 10:649-650

Hecker M. Wirksamkeit und Verträglichkeit von Efeuextrakt bei Patienten mit Atemwegserkrankungen. *Natura Med* 1999, 14(2):28-33

Hecker M, Runkel F, Völz A. Behandlung chronischer Bronchitis mit einem Spezialextrakt aus Efeublättern - multizentrische Anwendungsbeobachtung mit 1350 Patienten. *Forsch Kompl Klass Naturheil* 2002, 9:77-84

Hegener O, Prenner L, Runkel F, Baader SL, Kappler J, Häberlein H. Dynamics of β_2 -adrenergic receptor - ligand complexes on living cells. *Biochem* 2004, 43:6190-9

Hensel A, Deters AM, Müller G, Stark T, Wittschier N et al. Occurrence of N-phenylpropyl-L-amino acid amides in different herbal drugs and their influence on human keratinocytes, on human liver cells and on adhesion of *Helicobacter pylori* to the human stomach. *Planta Med* 2007, 73(2):142-150

Hiller K, Keipert M, Linzer B. Triterpensaponine. *Die Pharmazie* 1966, 21:713-750

Hocaoglu AB, Karaman O, Erge DO, Erbil G, Yilmaz O, Kivcak B et al. Effect of *Hedera helix* on lung histopathology in chronic asthma. *Iran J Allerg, Asthma Immunol* 2012, 11 (4):316-323

Hofmann D, Hecker M, Völp A. Efficacy of dry extract of ivy leaves in children with bronchial asthma – a review of randomized controlled trials. *Phytomedicine* 2003, 10:213-220

Holzinger F, Chenot JF. Systematic review of clinical trials assessing the effectiveness of ivy leaf (*Hedera helix*) for acute upper respiratory tract infections. *Evid Based Complement Alternat Med* 2011, 382789:1-9

Hong EH, Song JH, Sim A, Lee BR, Kwon BE, Song HH et al. Coadministration of *Hedera helix* L. Extract Enabled Mice to Overcome Insufficient Protection against Influenza A/PR/8 Virus Infection under Suboptimal Treatment with Oseltamivir. *Plos One* 2015, June 22:1-17

Hooshyar H, Talari S, and Feyzi F. Therapeutic effect of *Hedera helix* alcoholic extract against cutaneous leishmaniasis caused by *Leishmania major* in Balb/c mice. *Jundishj microbiol* 2014, 7(4):e9432

Hoppe HA. *Hedera*. Taschenbuch der Drogenkunde. De Gruyter Verlag, Berlin-New York 1981, 140

Hostettmann K. Saponins with molluscicidal activity from *Hedera helix* L. *Helvet Chim Acta* 1980, 63(3):606-09

Hostettmann K, Kizu H, Tomimori T. Molluscidial properties of various saponins. *Planta Med* 1982, 44(1):34-35

Ibrar M. Atomic absorption spectrophotometric analysis of the hypoglycemic trace elements of ivy (*Hedera helix* L.) leaves. *Hamdard Medicus* 2000, 43(3):68-71

Ibrar M, Ilahi I, Hussain F. The cytotoxic potential of ivy (*Hedera helix* L.) leaves. *Hamdard Medicus* 2001, 44(1):18-23

Ibrar M, Ilahi H, Hussain F. Hypoglycemic activity of *Hedera helix* L. leaves and the possible mechanism of action. *Pakist J Botan* 2003, 35(5):805-809

Ieven M, Vanden Berghe DA, Mertens F, Vlietinck A, Lammens E. Screening of higher plants for biological activities I. antimicrobial activity. *Planta Med* 1979, 36:311-21

Jahn E, Müller B. Efeublättertrockenextrakt. *Deutsch Apoth Zeit* 2000, 140(12):135-138

Jamal E, Gazala BH, Mabrouk E, Manal G, Abdulmotaleb Z, Abossa A et al. Cytotoxic activity of selected Libyan medicinal plants on human breast adenocarcinoma cell line (MCF-7). *Indon J Pharm* 2013, 24 (2):127-130

Jensen SR, Nielsen BJ, Dahlgren R (1975) Iridoid compounds, their occurrence and systematic importance in the Angiosperms. *Botaniska Notiser* 128:148–180

Jeong HG, Park HY. The prevention of carbon tetrachloride-induced hepatotoxicity in mice by α -hederin: inhibition of cytochrome P450 2E1 expression. *Biochem Mol Biol Intern* 1998, 45(1):163-170

Jeong HG. Suppression of constitutive and inducible cytochrome P450 gene expression by α -hederin in mice. *Biochem Molr Biol Intern* 1998, 46(5):1019-1026

Jøhnke H, Bjarnason B. Allergisk kontaktdermatitis over for efeu (*Hedera helix* L.). *Ugeskr Laeger* 1994, 156 (25):3778-3779

Jones JM, White IR, White JML, McFadden JP. Allergic contact dermatitis to English ivy (*Hedera helix*) - a case series. *Contact Derm* 2009, 60 (3):179-180

Julien J, Gasquet M, Maillard C, Balansard G, Timon-David P. Extracts of the ivy plant, *Hedera helix*, and their antihelminthic activity on liver flukes. *Planta Med* 1985, 51:205-208

Kardos P, Berck H, Fuchs K-H, Gillissen A, Klimek L, Morr H et al. Guidelines of the German respiratory society for diagnosis and treatment of adults suffering from acute or chronic cough. *Pneumologie* 2010, 64 (11):701-711

Kim SY, Son KH, Chang HW, Kang SS, Kim HP. Inhibition of mouse ear edema by steroid and triterpenoid saponins. *Archiv Pharm Res* 1999, 22(3):313-316

Kommission D Monographs (editors). *Hedera helix*. Bundesanzeiger Nr. 29a vom 12.02.1986 in der Fassung vom Bundesanzeiger Nr. 47 vom 08.03.1990

Kommission E Monographs (editors). *Hederae helicis folium* (Efeublätter). Bundesanzeiger Nr. 122 vom 06.07.1988

Kooperation Phytopharma. Stellungnahme zu "Efeu" – Kinderdosierungen vom 15.01.2003

Kraft K. Verträglichkeit von Efeublättertrockenextrakt im Kindesalter. *Zeitschr Phytother* 2004, 25:179-181

Kültür S. Medicinal plants used in Kirkclareli Province (Turkey). *J Ethnopharmacol* 2007, 111:341-364

Lässig W, Generlich H, Heydolph F, Paditz E. Wirksamkeit und Verträglichkeit efeuhaltiger Hustenmittel - Prospan® Kindersaft bei rezidivierenden obstruktiven Atemwegserkrankungen. *TW Pädiatrie* 1996, 9:489-491

Landgrebe H, Matusch R, Runkel F, Hecker M. Wirkung und Anwendung einer alten Heilpflanze. *Pharm Zeit* 1999, 144(35):11-15

Landgrebe H. Neue Wege zu wirksamen Inhaltsstoffen aus *Hedera helix* L. und deren Nachweis in Harn und Plasma. Dissertation, Philipps-Universität, Marburg 2002

Lang C, Röttger-Lüer P, Staiger C. A valuable option for the treatment of respiratory diseases: review on the clinical evidence of the ivy leaves dry extract EA 575®. *Planta Med* 2015a, 81:968-74

Lang C, Staiger C, Wegener T. Efeu in der pädiatrischen Praxis: Anwendung von EA 575® in der Therapie der akuten Bronchitis bei Schulkindern. [Ivy in everyday paediatric use: administration of EA 575® to schoolchildren for the treatment of acute bronchitis.] *Zeitschrift Phytother* 2015b, 36 (5):192-196

Lanza JP, Steinmetz MD, Pellegrin E, Mourgue M. Actions toxique et pharmacodynamique sur le rat d'extraits de lierre grimpant (*Hedera helix* L.). *Plantes méd phytothér* 1980, 14(4):221-229

Leonti M, Casu L, Raduner S, Cottiglia F, Floris C, Altmann KH et al. Falcarinol is a covalent cannabinoid CB1 receptor antagonist and induces pro-allergic effects in skin. *Biochem Pharmacol* 2010, 79 (12):1815-1826

Leskow P. Behandlung bronchialer Erkrankungen mit dem Phytotherapeutikum Prospan®. *Zeitschr Phytother* 1985, 6:61-64

- Liu J, Choudhuri S, Liu Y, Kreppel H, Andrews GK, Klaassen CD. Induction of metallothionein by α -hederin. *Toxicol Appl Pharmacol* 1993, 121:144-151
- Liu J, Liu Y, Bullock P, Klaassen CD. Suppression of liver cytochrome P450 by α -hederin: relevance to hepatoprotection. *Toxicol Appl Pharmacol* 1995, 134:124-131
- Liu YP, Liu J. Effect of α -hederin on hepatic detoxifying systems in mice. *Acta Pharm Sin* 1997, 18(1):33-36
- Liu B. Hederagenin from the leaves of ivy (*Hedera helix* L.) induces apoptosis in human LoVo colon cells through the mitochondrial pathway. *BMC Compl Altern Med* 2014, 14:412
- Loos M. Erfahrungen mit Prospan in der Klinik bei Krankheitsbildern des Respirationstraktes. *Mede Klin* 1958, 53(39):1693-1695
- Lorent J, Le Duff C S, Quetin-Leclercq J, Mingeot-Leclercq M. Induction of highly curved structures in relation to membrane permeabilization and budding by the triterpenoid saponins, α - and δ -hederin. *J biol chem* 2013, 288(20):14000-14017
- Lovell CR. *Hedera helix*. Plants and the skin. Blackwell Scientific Publications, Oxford 1993, 143-145
- Machado S, Silva E, Massa A. Occupational allergic contact dermatitis from falcarinol. *Contact Derm* 2002, 47:113-114
- Madaus G. *Hedera helix*. Lehrbuch der biologischen Heilmittel G. Thieme Verlag, Leipzig 1938, 1512-1517
- Mahillon V, Saussez S, Michel O. High incidence of sensitization to ornamental plants in allergic rhinitis. *Allergy* 2006, 61:1138-40
- Mahran GH, Hilal SH, El-Alfy TS. The isolation and characterisation of the emetine alkaloid from *Hedera helix*. *Planta Med* 1975, 27:127-132
- Maidannik V, Duk E, Kachalova O, Efanova A, Svojkina S, Sosnovskaja T. Efficacy of Prospan application in children's diseases of respiratory tract. *Ped, Toxicol Gynecol* 2003, 4:1-7
- Majester-Savornin B, Elias R, Diaz-Lanza AM, Balansard G, Gasquet M et al. Saponins of the ivy plant, *Hedera helix*, and their leishmanicidic activity. *Planta Med* 1991, 57:260-61
- Mansfeld HJ, Höhre H, Repges R, Dethlefsen U. Sekretolyse und Bronchospasmolyse. *TW Pädiatrie* 1997, 10(3):153-157
- Mansfeld HJ, Hohre H, Repges R, Dethlefsen U. Therapie des Asthma bronchiale mit Efeublätter-Trockenextrakt. *Münch Med Wochenschr* 1998, 140(3):26-30
- März RW, Matthys H. Phytomedicines in the treatment of diseases of the lower respiratory tract. What is proven? In: Loew D, Rietbrock N, editors. *Phytopharmaka III - Forschung und klinische Anwendung*. Steinkopff, Darmstadt 1997, 161-178
- Mayer H, Pfandl A, Grigorieff A, Zickner J. Der Efeu - eine alte Kult- und Heilpflanze. *Pharm Zeit* 1987, 132:2673-2676
- Mayer JG. Efeu - *Hedera helix* L.: Kulturhistorisches Porträt einer Arzneipflanze. Ivy (*Hedera helix*). Cultural and historical portrait of a medicinal plant. *Zeitschr Phytother* 2010, 31 (5):271-274
- Mba Gachou C, Laget M, Guiraud-Dauriac H, De Meo M, Elias R, Duménil G. The protective activity of α -hederine against H_2O_2 genotoxicity in HepG2 cells by alkaline comet assay. *Mut Res* 1999, 445:9-20

Mendel M, Chlopecka M, Dziekan N and Wiecheteck M. The effect of the whole extract of common ivy (*Hedera helix*) leaves and selected active substances on the motoric activity of rat isolated stomach strips. *J Ethnopharmacol* 2011, 134: 796-802

Mendel M, Chlopecka M, Dziekan N, Karlik and W, Wiecheteck M. Participation of cholinergic pathways in α -hederin-induced contraction of rat isolated stomach strips. *Phytomedicine* 2012, 19: 591-595

Mendel M, Chlopecka M, Dziekan N, Karlik W and Wiecheteck M. Participation of extracellular calcium in α -hederin-induced contractions of rat isolated stomach strips. *J Ethnopharmacol* 2013, 146: 423-426

Meincke R, Pokladnikova J, Niedrig DF, Russmann S, Meyboom RHB. Acute hypersensitivity reactions to herbal remedies in children reported to the WHO VigiBase® database. Conference Paper in *Drug Safety* October 2015, 38: 954

Metcalfe DJ. *Hedera helix* L. *J Ecol* 2005, 93: 632-648

Meyer-Wegener J, Liebscher K, Hettich M, Kaster HG. Efeu versus Ambroxol bei chronischer Bronchitis. *Zeitschr Allgemeinmed* 1993, 69(3): 61-66

Mills S, Bone K. Saponins. In: Principles and practice of phytotherapy. Churchill Livingstone, Edinburgh 2000, 43-47

Mitchell JC, Rook AJ. Hedera. Botanical Dermatology. Greengrass, Vancouver 1979, 121-123

Mitchell JC. Allergic contact dermatitis from *Hedera helix* and *Brassica actinophylla*. *Contact Derm* 1981, 7(3): 158-159

Morfin-Maciel BM, Rosas-Alvarado A, Velázquez-Sámano G. Anaphylaxis due to ingestion of ivy syrup (*Hedera helix*). Report of two cases. *Rev Alerg Mex* 2012, 59: 31-6 [Spanish]

Morce AH, McGarvey L, Pavord I. BTS Guidelines: Recommendations for the management of cough in adults. *Thorax* 2006, 61: 1-24

Moulin-Traffort J, Favel A, Elias R, Regli P. Study of the action of α -hederin on the ultrastructure of *Candida albicans*. *Mycoses* 1998, 41(9-10): 411-416

Mühlendahl KE, Oberdissen U, Bunjes R, Ritter S, editors. Vergiftungen im Kindesalter, Ferdinand Enke Verlag, Stuttgart 1995, 347

Müller B, Bracher A. Wirksamkeit und Verträglichkeit von Efeublättertrockenextrakt: Ergebnisse einer erweiterten Anwendungsbeobachtung bei 372 Kindern im Alter von 0-16 Jahren. In: Schulz V, Rietbrock N, Roots I, Loew D, editors. Phytopharmaka VII - Forschung und klinische Anwendung. Steinkopff, Darmstadt 2002, 55-58

Müller-Jakic B. Current scientific information concerning "Hederae helicis folium" (ivy leafs) since 1987 – taking earlier relevant studies into consideration (drafted in accordance with the "Notice to Applicants") Kooperation Phytopharmaka GbR 1998, 1-25

Mulkijanyan K, Novikova Zh, Sulakvelidze M, Getia M, Mshvildadze V, Dekanosidze G. Ivy water extracts as gastric ulcer preventive agents. *Georgian med news* 2013, 224: 63-6

Murdoch SR, Dempster J. Allergic contact dermatitis from carrot. *Contact Dermatitis* 2000, 42(4): 236

Olariu L, Rosoiu N, Constantinovici M, Drumea V. Evolution of B16 melanoma following treatment with several bioactive plant compounds. *FEBS Journal* 2007, 274(1): 166

Orhan DD, Özceli K B, Hosbas S, Vural M. Assessment of antioxidant, antibacterial, antimycobacterial, and antifungal activities of some plants used as folk remedies in Turkey against dermatophytes and yeast-like fungi. *Turkish J Biol* 2012, 36 (6):672-686

Ortiz de Urbina JJ, Martin ML, Sevilla MA, Montero MJ, Carron R, San Roman L. Antispasmodic activity on rat smooth muscle of polyphenol compounds caffeic and protocathechic acids. *Phytother Res* 1990, 4(2): 71-76

Özdemir C, Schneider LA, Hinrichs R, Staib G, Weber L, Weiss JM et al. Allergische Kontaktdermatitis auf Efeu (*Hedera helix* L.). *Der Hautarzt* 2003, 54(10):966-969

Paulsen E, Christensen LP, Andersen KE. Dermatitis from common ivy (*Hedera helix* L. subsp. *helix*) in Europe: past, present, and future. *Cont Derm* 2010, 62:201-9

Pharmacopée Francaise VIII edition. Tisanes. 1965:1224-1226

Pokladnikova J, Meyboom RHB, Meincke R, Niedrig D, Russmann S. Allergy-like immediate reactions with herbal medicines: a retrospective study using data from VigiBase®. *Drug Safety* 2016, 39(5):455-64.

Polizzi A, Incopora G, Ruggieri M. Dystonia as acute adverse reaction to cough suppressant in a 3-year-old girl. *Eur J Ped Neurol* 2001, 5(4):167-8

Quentin-Leclercq J, Elias R, Balansard G, Bassleer R, Angenot L. Cytotoxic activity of some triterpenoid saponins. *Planta Med* 1992, 58:279-281

Rai A. The effects of alcoholic *Hedera helix* plant extract on ehrlich ascitic tumour cells *in vitro* and its antitumour activities *in vivo*. *Intern J Pharm Sci Rev Res* 2011, 11 (2):47-50

Rai A. The anti-inflammatory and antiarthritic properties of ethanol extract of *Hedera helix*. *Indian J Pharm Sci* 2013, 75:99-102

Rao GS, Sinsheimer JE, Cochran KW. Antiviral activity of triterpenoid saponins containing acylated β-amyrin aglycones. *J Pharm Sci* 1974, 63:471-473

Rath F. Klinische Prüfung der Wirksamkeit des Hustenmittels Prospan. *Fortschr There* 1968, 22:1015-1016

Ridoux O, Di Giorgi C, Delmas F, Elias R, Mshvildadze V et al. *In vitro* antileishmanial activity of three saponins isolated from Ivy, α-Hederin, β-Hederin and Hederacolchiside A1, in association with pentamidine and amphotericin B. *Phytother Res* 2001, 15(4):298-301

Roed - Petersen J. Allergic contact hypersensitivity to ivy (*Hedera helix*). *Cont Derm* 1975, 1:57

Roth R. Efeublätter wirken sekretolytisch und bronchospamolytisch - Anwendungsbeobachtung bestätigt Wirksamkeit der Behandlung mit Efeublätter-Trockenextrakt. *PädiatrNachr* 2000, Sonderdruck: 1-3

Rudkowski Z, Latos T. Inhalationsbehandlung chronischer Bronchitiden im Kindesalter mit Prospan. *Ärztliche Praxis* 1979, 31(10):342-346

Runkel F, Prenner L, Häberlein H. Ein Beitrag zum Wirkmechanismus von Efeu. *Pharm Zeit* 2005, 150(4):19-25

Sánchez-Pérez J, Córdoba S, Hausen BM, de Vega MJ, Aragüés M, García-Díez A. Allergic contact dermatitis from common ivy confirmed with stored allergens. *Contact Dermatitis* 1998, 39(5):259-260

Schaefer A, Kehr MS, Giannetti BM, Bulitta M, Staiger C. A randomized, controlled, double-blind, multi-center trial to evaluate the efficacy and safety of a liquid ivy leaves dry extract (EA 575) vs. placebo in the treatment of adults with acute cough. *Pharm* 2016, 71 (9):504-509

Schapowal A. Phytopharmaka bei Atemwegsinfekten in der Pädiatrie. *Zeitschr Phytother* 2007, 28(4):174-180

Schmidt M, Thomsen M, Schmidt U. Suitability of ivy extract for the treatment of paediatric cough. *PhytotherRes* 2012, 26 (12) 1942-7 (wileyonlinelibrary.com) DOI:10.1002/ptr.4671

Schmidt O. Biopharmazeutische Charakterisierung von Efeublättertrockenextrakten durch *in vivo* und *in vitro* Studien. Dissertation, Ruprecht-Karls-Universität, Heidelberg 2003, 109-210

Schottk W. Untersuchungen über die Eignung von *Hedera helix*-Präparaten für die Prophylaxe bzw. Therapie von Reizgasvergiftungen. *Berufsderm* 1972, 20:88-92

Shi JZ, Liu GT. Effect of α -hederin and sapindoside B on hepatic microsomal cytochrome P-450 in mice. *Acta Pharm Sin* 1996, 17(3):264-266

Sieben A, Prenner L, Sorkalla T, Wolf A, Jakobs D, Runkel F, Häberlein H. α -Hederin, but not hederacoside C and hederagenin from *Hedera helix*, affects the binding behavior, dynamics, and regulation of beta 2-adrenergic receptors. *Biochem* 2009, 48(15):3477-3482

Song JH, Yeo SG, Hong EH, Lee BR, Kim JW, Kim JH et al. Antiviral activity of hederasaponin B from *Hedera helix* against enterovirus 71 subgenotypes C3 and C4a. *Biomol ther* 2014, 22(1):41-46

Song KJ, Shin YJ, Lee KR, Lee EJ, Suh YS, Kim KS. Expectorant and antitussive effect of *Hedera helix* and *Rhizoma coptidis* extracts mixture. *Yonsei Med J*, 2015, 56 (3):819-824

Stauss-Grabo M. Untersuchungen zur Pharmakokinetik des Efeu-Saponins α -Hederin in Ratten. PhD Thesis, Philipps University Marburg 2008

Stauss-Grabo M, Häberlein H, Runkel F. EFEU - Ein modernes Phytopharmakon. *Pharmazie in Unserer Zeit* 2008, 37(6):484-485

Stauss-Grabo M, Atiye S, Warnke A, Wedemeyer RS, Donath F, Blume HH. Observational study on the tolerability and safety of film-coated tablets containing ivy extract (Prospan® Cough Tablets) in the treatment of colds accompanied by coughing. *Phytomedicine* 2011, 18:433-6

Steinmetz EF. Hederae helicis folia. *Pharm Biol* 1961, 1(3):93-95

Stöcklin P. Klinische Erfahrungen mit dem Hustenmittel Prospan. *Praxis* 1959, 41:934-938

Süleyman H, Mshvildadze V, Gepdiremen A, Elias R. Acute and chronic antiinflammatory profile of the ivy plant, *Hedera helix*, in rats. *Phytomed* 2003, 10(5):370-374

Sweetmann SC. Ivy. Martindale: The Complete Drug Reference, 35th ed, Pharmaceutical Press 2007, 2111

Tabali M, Ostermann T, Jeschke E, Witt CM, Matthes H. Adverse drug reactions for CAM and conventional drugs detected in a network of physicians certified to prescribe CAM drugs. *J mancare pharm* 2012, 18 (6):427-38

Teat DW, Ellis MD. Toxic indoor plants. *US Pharmacist* 1981, 6:38-39

Tedlaouti F, Gasquet M, Delmas F, Timon-David P, Elias R et al. Antitrypanosomal activity of some saponins from *Calendula arvensis*, *Hedera helix* and *Sapindus mukorossi*. *Planta Med* 1991, 57:A78

Thormann H, Paulsen E. Contact urticaria to common ivy (*Hedera helix* cv. 'Hester') with concomitant immediate sensitivity to the labiate family (Lamiaceae) in a Danish gardener. *Contact Derm* 2008, 59(3):179-80

Timon-David P, Julien J, Gasquet M, Balansard G, Bernard P. Recherche d'une activité antifongique de plusieurs principes actifs extraits du hêtre grimpant. *Ann Pharm Fr* 1980, 38(6):545-552

Trute A, Gross J, Mutschler E, Nahrstedt A. *In vitro* antispasmodic compounds of the dry extract obtained from *Hedera helix*. *Planta Med* 1997, 63(2):125-129

Turton PHJ. Poisoning by ivy. *Br Med J* 1925, 3372:294

Uddin G, Rauf A, Qaisar M, Latif A, Ali M. Preliminary phytochemical screening and antimicrobial activity of *Hedera helix* L., Middle East, *J Sci Res* 2011, 8 (1):198-202

Unkauf M, Friedrich M. Bronchitis bei Kindern - Klinische Studie mit Efeublätter-Trockenextrakt. *Der Bayr Intern* 2000, 4:1-4

Van den Bout-van den Beukel CJP, Koopmans PP, van der Ven AJAM, De Smet PAGM, Burger DM. Possible drug-metabolism interactions of medicinal herbs with antiretroviral agents. *Drug Metabol Review* 2006, 38:477-514

Valnet J. Lierre grimpant. In: Maloine, editor. *Phytothérapie*, 5th ed. Paris 1983, 506-509

Vasconcelos Pinela AF, Cunha D, Maio P, Amaro C, Santos R, Inácio F. Allergic phytodermatitis to common ivy (*Hedera helix* spp.) - A case report. *Allergy: Eur J Allerg Clin Immun* 2010, 65 :596

Vila R, Cañigueral S. Ivy leaf in the treatment of respiratory conditions: preclinical and clinical evidences. *RevFitoter* 2011, 11(1) :5-20

Villani P, Orsiere T, Sari-Minodier I, Bouvenot G, Botta A. Etude *in vitro* de l'activité antimutagène de l'alphahéderine. *Ann Biol Clin* 2001, 59(3):339-353

Vogel G, Marek ML. Zur Pharmakologie einiger Saponine. *Arzneimitt.-Forsch* 1962, 12(8):815-825

Vogel G. Zur Pharmakologie von Saponinen. *Planta Med* 1963, 11(3):362-376

Vogelmeier C, Buhl R, Criée CP, Gillissen A, Kardos P, Köhler D et al. Leitlinie der Deutschen Atemwegsliga und der Deutschen Gesellschaft für Pneumologie und Beatmungsmedizin zur Diagnostik und Therapie von Patienten mit chronisch obstruktiver Bronchitis und Lungenemphysem (COPD). [Guidelines for the diagnosis and therapy of COPD issued by Deutsche Atemwegsliga and Deutsche Gesellschaft für Pneumologie und Beatmungsmedizin.] *Pneumologie* 2007, 61:1-40

Wagner H, Reger H. Folium Hedera-Extrakte - HPLC Analyse. *Deutsch Apoth Zeit* 1986, 126(48):2613-2617

Wagner H, Wiesnauer M. Saponin-Drogen. In: *Phytotherapie*. Fischer Verlag, Stuttgart 1995, 100-108

WHO. GOLD guideline. Global initiative for chronic obstructive lung disease, 2006, 1-100

Wichtl M. Teedrogen und Phytopharmaka. 2nd ed. Wissenschaftliche Verlagsgesellschaft mbH, Stuttgart 1989, 139-141

Wichtl M, Anton R. *Hederae folium*. Plantes Thérapeutiques. Tradition, pratique officinale, science et thérapeutique. Ed. Tec. et Doc., Paris 1999, 251-253

Wichtl M (editor). *Hedera folium. Herbal drug and Phytopharmaceuticals* 3rd ed. MedPharm GmbH Scientific Publishers, Marburg 2004, 274-277

Williamson EM. Ivy. Potter's Herbal Cyclopaedia. Limited 2003, 237-238

Wolters B. Die Verbreitung antibiotischer Eigenschaften bei Saponindrogen. *Deutsch Apoth Zeit* 1966, 106:1729-1733

Wulff G. Neuere Entwicklungen auf dem Saponingebiet. *Deutsch Apoth Zeit* 1968, 108(23):797-808

Yesudian PD, Franks A. Contact dermatitis from *Hedera helix* in a husband and wife. *Contact Derm* 2002, 46:125-126

Zeil S, Schwanbeck U, Vogelberg C. Tolerance and effect of an add-on treatment with a cough medicine containing ivy leaves dry extract on lung function in children with bronchial asthma. *Phytomedicine* 2014, 21:1216-20

References assessed, but not cited in the assessment report:

- Babadjamian A, Elias R, Faure R, Vidal-Ollivier E, Balansard G. Two-dimensional NMR studies of triterpenoid glycosides. ¹H and ¹³C NMR assignments of herderasaponin C [3-O- α -L-rhamnopyranosyl-(1 \rightarrow 2)- α -L-arabinopyranosyl-hederagenin 28-O- α -L-rhamnopyranosyl-(4)- β -D-glucopyranosyl-(1 \rightarrow 6)- β -D-glucopyranosylester]. *Spectr Letts* 1988, 21(7):565-573
- Bader G. Pharmakologische und biopharmazeutische Bewertung von Triterpensaponinen. *Pharmazie* 1994, 49:391-400
- Bedir E, Kirmizipekmez H, Calis I, Sticher O. Two new hederagenin glycosides from the fruits of *Hedera helix* L. 46. Kongress der Gesellschaft für Arzneipflanzenforschung "Quality of Medicinal Plants and Herbal Medicinal Products" in Vienna. 1998, 58
- Bedir E, Kirmizipekmez H, Sticher O, Calis I. Triterpene saponins from the fruits of *Hedera helix*. *Phytochemistry* 2000, 53(8): 905-909
- Berger F. Handbuch der Drogenkunde Bd. 2, Wilhelm Maudrich Verlag, Wien 1950, 156-158
- Büechi S, Vögelin R, von Eiff MM, Ramos M, Melzer J. Open trial to assess aspects of safety and efficacy of a combined herbal cough syrup with ivy and thyme. *Forsch Komplem Klass Natur* 2005, 12(6):328-332
- Bühring U. Efeu - *Hedera helix* L. *Naturheilpraxis* 2003, 05:669-671
- Busser C. Baies, fruits et pseudo-fruits toxiques utilisés en médecine populaire ou en phytothérapie. *Phytothérapie* 2007, 5(1):31-36
- Calnan CD. Dermatitis from Ivy (*Hedera canariensis variegata*). *Contact Dermatitis* 1981, 7(2):124
- Campbell A. Poisoning in small animals from commonly ingested plants. *In Practice* 1998, 20(10):587-591
- Cortesi R, Weber HR. Recherches de quelques vitamines dans le lierre grimpant. (*Hedera helix*. L.). *Boll Chim Farm* 1958, 97:131-137
- de Costa F, Yendo ACA, Fleck JD, Gosmann G, Fett-Neto AG. Immunoadjuvant and anti-inflammatory plant saponins: Characteristics and biotechnological approaches towards sustainable production. *Mini-Rev Medl Chem* 2011, 11 (10):857-880
- De Smet PAGM. Herbal medicine in Europe - relaxing regulatory standards. *New England J Med* 2005, 352(12):1176-1178
- Dorsch W. Phytopharmaka in der pädiatrischen Praxis 5. Phytotherapiekongress Bonn 5.-6.11.1993
Abstract: 6-8
- Dorsey CS. Algerian ivy dermatitis – A California disease. *California Medicine* 1959, 90(2):155-159
- Elias R, Diaz Lanza AM, Vidal-Ollivier E, Balansard G, Faure R, Babadjamian A. Triterpenoid saponins from the leaves of *Hedera helix*. *J Nat Prod* 1991, 54(1): 98-103
- Eller I, Lode H. Pathophysiologie des Hustens. In: Tyrrell DAJ, editor. Erkältungskrankheiten - ein Lehrbuch für die Praxis. Gustav Fischer Verlag, Stuttgart, Jena, New York 1996, 163-173
- Fahay F, Stocks N, Thomas T. Quantitative systematic review of randomised controlled trials comparing antibiotic with placebo for acute cough in adults. *Br Med J* 1998, 316:906-910
- Forth W. Pflanzliche Bronchospasmolytika und Expektorantien. *Therapiewoche* 1984, 34:4089-4096

Francis G, Kerem Z, Makkar HP, Becker K. The biological action of saponins in animal systems: a review. *Br J Nutr* 2002, 88(6):587-605

Fröhling D. Saponin - Bestimmung in Hedera Extrakten. *Pharmazie* 1969, 24:704

Gafner F, Reynolds G, Rodriguez E. The diacetylene 11,12-dehydrofalcarkinol from *Hedera helix*. *Phytochemistry* 1989, 28:1256-1257

Gillissen A, Tasci S, Ewig S, Schäfer H, Zielen S. Sinn und Unsinn von Antitussiva. *Internist* 2001, 42:134-145

Glasl H, Ihrig M. Quantitative Bestimmung von Triterpensaponinen in Drogen. *Pharm Zeit* 1984, 129:2619-2622

Grosbois M. Biosynthese des acides gras au cours du développement du fruit et de la graine du lierre. *Phytochem* 1971, 10:1261-1273

Hahn G, Hahn M. Der Efeu, eine alte Kultur- und Medizinalpflanze. *Östereich Apotheker Zeit* 1982, 52:954-959

Hahn HL. Husten - Mechanismen, Pathophysiologie und Therapie. *Deutsch Apotheker Zeit* 1987, 127(5):3-26

Hambly EM, Wilkinson DS. Sensitivity to variegated Ivy (*Hedera canariensis*). *Contact Derm* 1978, 4:239-240

Hansen NC, Skriver A, Brorsen-Riis L, Balslev S, Evald T, Maltbaek N et al. Orally administered N-acetylcysteine may improve general well-being in patients with mild chronic bronchitis. *Respir Med* 1994, 88(7):531-535

Hansen L, Hammershøi O, Boll PM. Allergic contact dermatitis from falcarinol isolated from *Schefflera arboricola*. *Contact Derm* 1986, 14:91-93

Hegi G (editor). *Hedera*. In: *Illustrierte Flora von Mitteleuropa* Bd. V, Teil 2, Verlag Paul Parey, Berlin 1975, 914-925

Hillman JR, Knights BA, McKail R. Sterols of the adult and juvenile forms of ivy, *Hedera helix*. *Lipids* 1975, 10(9):542-544

Hoppe JE. Keuchhusten bei Erwachsenen keineswegs selten - jedoch oft verkannt. *Intern Praxis* 1999, 39:53-57

Imhof E, Russi E, Perruchoud AP. Pharmakotherapie des Hustens. *Schweizer Med Wochenschr* 1988, 118(29):1067-1072

Iszard MB, Liu J, Klaassen CD. Effect of several metallothionein inducers on oxidative stress defense mechanisms in rats. *Toxicol* 1995, 104:25-33

Jensen SR, Nielsen BJ, Dahlgren R. Iridoid compounds, their occurrence and systemic importance in the angiosperms. *Botaniska Notiser* 1975, 128:148-180

Jungmayr P. Phytotherapie: Efeuextrakte bei Husten. *Deutsche Apotheker Zeit* 2007, 147(46):60-61

Jury SL. The Euro+Med treatment of *Hedera* (Araliaceae) - recent studies and a new name. *Willdenowia* 2004, 34(2):351-352

Kartnig T, Wegschaider O, Ri CY. Ein Beitrag zur Bestimmung von Saponinen in Drogen. Determination of saponins in medicinal plants. *Planta Med* 1972, 21:29-34

Kartnig T, Ri CY, Wegschaider O. Untersuchungen über die Aufbereitung und die Stabilität von Saponindrogen. *Planta Med* 1972, 22:127-135

Khan JA, Siddiqui MA, Nikhat S. Contraceptives in Greek literature: A review. *Intern Res J Pharm* 2013, 4(9):22-24

Kemmerich B, Eberhardt R, Stammer H. Efficacy and tolerability of a fluid extract combination of Thyme herb and Ivy leaves and matched placebo in adults suffering from acute bronchitis with productive cough. *Arzneimittelforsch* 2006, 56(9):652-660

König P. Therapie einer primär-biliären Zirrhose mit *Hedera helix*. *AHZ* 1991, 236(4):143-149

Kraft K. Symptomatische Phytotherapie bei Husten: Stellenwert pflanzlicher Antitussiva und Expektorantien. (Symptomatic phytotherapy in coughs: Importance of plant antitussives and expectorants). *Pharmazie in Unserer Zeit* 2008, 37(6):478-483

Krakauer G, Rattenhuber S. Therapeutical experiences with Prospan® for the aerosol application in horses. *Der Praktische Tierarzt* 1984, 11:980-986

Lacaille-Dubois MA. Biologically and pharmacologically active saponins from plants: recent advances. In: Marston A and Oleszek W, (editors. *Saponins in Food, Feedstuffs and Medicinal Plants*. Kluwer Academic Publisher, Dordrecht 2000, 205-218

Lawrence GHM, Schulze AE. The cultivated Hederas. *Gentes Herbarium* 1942, 6:107-173

Lawrence GHM. The cultivated ivies. *Morris Arboretum Bull* 1965, 7:19-31

Liebscher K. Efeu versus Ambroxol bei chronischer Bronchitis. 5. Phytotherapietkongress Bonn 1993, Abstract band 42

Loew D, Schrödter A, Schilcher H. Phytopharmaka bei katarrhalischen Erkrankungen der oberen und unteren Atemwege. In: Loew D, Rietbrock N, editors. *Phytopharmaka III - Forschung und klinische Anwendung*, Steinkopff, Darmstadt 1997, 179-190

Mahe-Quino M, Rossinyol G, Foucaud A. Empoisonnement mortel de poulets par le graines de lierre. *Plant Méd Phytothér Tome IX* 1975, 3:182-186

Mansouri L, Henry M. An ontogenetic study on the sapogenin contents of ivy. *Pharmazie* 1995, 50:642-643

Marzian O. Behandlung der akuten Bronchitis bei Kindern und Jugendlichen. Anwendungsbeobachtung bestätigt Nutzen und Unbedenklichkeit einer Flüssigextrakt-Kombination aus Thymiankraut und Efeublättern. *MMW, Fortschr Med. Originalia* 2007, 149(2):69-74

Mayer A, Hahn G. Der Efeu. *Natur & Heilen* 1986, 63:366-367

Menßen HG. Gewöhnlicher Efeu. In: Menßen HG, editor. *Phytotherapeutische Welt*, PMI Verlag GmbH, Frankfurt am Main 1983, 96-99

Meyer EA. Der Efeu. *Pharm Rund* 1994, 4:33

Mutschler E. Arzneimittelwirkungen. 8th ed. Wissenschaftliche Verlagsgesellschaft, Stuttgart 2001, 621

Pahlow M. Das große Buch der Heilpflanzen. Gräfe und Unzer Verlag, München 1979, 111-112

Plohmann B, Bader G, Streich S, Hiller K, Franz G. Immunomodulatory effects of triterpenoid saponins. *Eur J Pharm Sci* 1994, 2(1-2):120

Rao GS, Sinsheimer JE, Cochran KW. Antiviral activity of triterpenoid saponins containing acylated β -amyrin aglycones. *J Pharm Sci* 1974, 63(3):471-473

Raynaud J. Les heterosides flavononiques d'Hedera helix. *Plantes Méd Phytothér* 1982, 16:318-320

Reuter HD. Behandlung von Erkältungskrankheiten mit Phytopharmaka. *Sozialpädiatrie* 1992, 14/3:194-203

Reuter HD, editor. Therapie mit Phytopharmaka. Pharmakologie, Indikationen, Dosierungen. Gustav Fischer Verlag, Stuttgart 1997, 61-62

Rocha LG, Almeida JRGS, Macêdo RO, Barbosa-Filho JM. A review of natural products with antileishmanial activity. *Phytomed* 2005, 12:514-535

Roth L, Daunderer M, Kormann K. Giftpflanzen Pflanzengifte. 4th ed. Nikoverlag, Hamburg 2006, 390-391

Rudkowski Z, Latos T. Erfahrungsbericht. *Hedera helix*: wirksam bei Bronchitis im Kindesalter. *Ärztliche Praxis* 1980, 80:2561-2662

Sant'Ambrogio G, Tsubone H, Sant'Ambrogio FB. Sensory information from the upper airway: role in the control of breathing. *Respiration Physiology* 1995, 102:1-16

Schilcher H. Phytotherapie in der Kinderheilkunde: Handbuch für Ärzte und Apotheker. 3rd ed. Wissenschaftliche Verlagsgesellschaft mbH, Stuttgart 1999, 36-41

Schlenger R. Wirkmechanismus von Efeuextrakt entschlüsselt. *Deutsch Apoth Zeit* 2003, 143:54-55

Schlösser E. Die Rolle von Saponinen im Resistenzmechanismus gegen Pilze. *Pflanzen Krankheiten* 1973, 11/12:704-710

Schmidt M. Mit Efeu gegen den Husten. *PTA Heute* 2003, 11:42-44

Schulz V, Hänsel R. Rationale Phytotherapie. Springer Berlin, Heidelberg, New York 1996, 167-168

Segal R, Schlosser E. Role of glycosidases in the membranolytic, antifungal action of Saponins. *Archiv Microbiol* 1975, 104:147-150

Sidhu GS, Oakenfull DG. A mechanism for the hypocholesterolaemic activity of saponins. *Brit J Nutr* 1986, 3:643-649

Thoms H, editor. Handbuch der praktischen und wissenschaftlichen Pharmazie. Vol. 2. Verlag Urban und Schwarzenberg, Berlin Wien 1938, 1370

Tobler F. Die Gattung Hedera. Gustav Fischer Verlag, Jena 1912, 1-151

Tronchet J. Physiologie vegetale. *CR Acad Sci Paris* 1964, 258:2390-2392

Trute A, Nahrstedt A. Identification and quantitative analysis of phenolic compounds from the dry extract of *Hedera helix*. *Planta Med* 1997, 63:177-179

Tschesche R, Schmidt W, Wulff G. Reindarstellung und Strukturermittlung der Saponine des Efeus (*Hedera helix* L.). *Zeitschr Naturforsch* 1965, 20b:708-709

Urban R. Physiologische Untersuchungen über einige Flavonoide und Oxyzimtsäuren. *Planta* 1958, 52:47-64

Wagner L, Cramer H, Klose P, Lauche R, Gass F, Dobos G, Langhorst J. Herbal Medicine for Cough: a Systematic Review and Meta-Analysis. *ForschKomplem* 2006, 22(6):359-68

Walther U, Dittrich K, Gelbrich G, Schöpke T. Effects of saponins on the water solubility of different model compounds. *Planta Med* 2001, 67:49-54

Wemmer U. Expektoranzien und Antitussiva bei Husten. *AJ* 1996, 18(10):12-22

Young I. Steroidal compounds in *Phaseolus* and *Hedera*. *Trans. Botan Soc Edinburgh* 1976, 42(4):533