



24 November 2021  
EMA/HMPC/625624/2015  
Committee on Herbal Medicinal Products (HMPC)

## List of references supporting the assessment of *Saccharomyces cerevisiae CBS 5926* Final

**The European Medicines Agency acknowledges that copies of the underlying works used to produce this monograph were provided for research only with exclusion of any commercial purpose.**

Adam J, Barret A, Barret-Bellet C. Essais cliniques contrôlés en double insu de l'ultra-levure lyophilisée. Etude multicentrique par 25 médecins de 388 cas. *Gaz Med Fr* 1977, 84:2072-2078 [in French]

Akhondi-Meybodi M, Rahimian M, Salmanroghani H, Amirbeigy M, Baghbanian M, Ghelmani SY. Study of the effect of probiotic *Saccharomyces boulardii* on the treatment of irritable bowel syndrome. *J Biol Today's World* 2014, 3(7):152-156

Akil I, Yilmaz O, Kurutepe S, Degerli K, Kavukcu S. Influence of oral intake of *Saccharomyces boulardii* on *Escherichia coli* in enteric flora. *Pediatr Nephrol* 2006, 21:807-810

Amorissani Folquet M, Saizonou R, Guedehoussou T, Vandenplas Y. *Saccharomyces boulardii* in acute gastroenteritis in children. *Webmed Central Paediatrics* 2011, 2(11):WMC001627

Asmat S, Shaukat F, Asmat R, Bkhhat H, Asmat T. Clinical efficacy comparison of *Saccharomyces boulardii* and lactic acid as probiotics in acute pediatric diarrhea. *JCPSP* 2018, 28(3):214-217

Bafutto M, de Almeida JR, Leite NV, Costa M, de Oliveira EC, Resende-Filho J. Treatment of diarrhea-predominant irritable bowel syndrome with mesalazine and/or *Saccharomyces boulardii*. *Arq Gastroenterol* 2013, 50(4):304-309

Barc MC, Charrin-Sarnel C, Rochet V, Bourlioux F, Sandré C, Boureau H, et al. Molecular analysis of the digestive microbiota in a gnotobiotic mouse model during antibiotic treatment: influence of *Saccharomyces boulardii*. *Anaerobe* 2008, 14(4):229-233

Bartlett JG. Antibiotic-associated diarrhea. *N Engl J Med* 2002, 346:334-339

Bassetti S, Frei R, Zimmerli W. Fungemia with *Saccharomyces cerevisiae* after treatment with *Saccharomyces boulardii*. *Am J Med* 1998, 105(1):71-72

Bedi MK, Shenefelt PD. Herbal therapy in dermatology. *Arch Dermatol* 2002, 138:232-242



Benhamou PH, Berlier P, Danjou G, Plique O, Jessuel D, DuPont C. La diarrhée aux antibiotiques chez l'enfant: une étude informatisée en double aveugle chez des patients ambulatoires comparant un agent protecteur et un agent probiotique. *Méd Chir Dig* 1999, 28:163-168 [in French]

Besirbellioglu BA, Ulcay A, Can M, Erdem H, Tanyuksel M, Avci IY, et al. *Saccharomyces boulardii* and infection due to *Gardia lamblia*. *Scand J Infect Dis* 2006, 38:479-481

Billoo AG, Memon MA, Khaskhelli SA, Murtaza G, Iqbal K, Saeed Shekhani M, et al. Role of a probiotic (*Saccharomyces boulardii*) in management and prevention of diarrhoea. *World J Gastroenterol* 2006, 28(12):4557-4560

Bin Z, Ya-Zheng X, Zhao-Hui D, Bo C, Li-Rong J, Vandenplas Y. The efficacy of *Saccharomyces boulardii* CNCM I-745 in addition to standard *Helicobacter pylori* eradication treatment in children. *Pediatr Gastroenterol Hepatol Nutr* 2015, 18(1):17-22

Biocodex. Compositions and methods for preventing infections. Patent WO 2015189337 A1. 17.12.2015; Biocodex homepage. Available at: <https://www.biocodex.com/en/therapeutic-areas-products/>

Blackwell B, Marley E. Interactions of yeast extracts and their constituents with monoamine in man and rat. *Brit J Pharmacol* 1966; 26:142-161

Blaschek W, Hilgenfeldt U, Holzgrabe U, Reichling J, Ruth P, Schulz. HagerROM. Hagers Enzyklopädie der Arzneistoffe und Stoffe. Springer Verlag, Berlin Heidelberg 2013 [in German]

Blehaut H, Massot J, Elmer GW, Levy RH. Disposition kinetics of *Saccharomyces boulardii* in man and rat. *Biopharm Drug Dispos* 1989, 10:353-364

Bleichner G, Bléhault H, Mentec H, Moyse D. *Saccharomyces boulardii* prevents diarrhea in critically ill tube-fed patients. A multicenter, randomized, double-blind placebo-controlled trial. *Intensive Care Med* 1997, 23:517-523

Bravo MV, Bunout D, Leiva L, de la Maza MP, Barrera G, de la Maza J et al. Effect of probiotic *Saccharomyces boulardii* on prevention of antibiotic-associated diarrhea in adult outpatients with amoxicillin treatment. *Rev Méd Chile* 2008, 136:981-988 [in Spanish]

Burande MA. Comparison of efficacy of *Saccharomyces boulardii* strain in the treatment of acute diarrhea in children: a prospective, single-blind, randomized controlled clinical trial. *J Pharmacol Pharmacother* 2013, 4(3):205-208

Buts JP, Bernasconi P, van Craynest MP, Maldague P, de Meyer R. Response of human and rat small intestinal mucosa to oral administration of *Saccharomyces boulardii*. *Pediatr Res* 1986, 20:192-196

Buts JP, Bernasconi P, Vaerman J-P, Dive C. Stimulation of secretory IgA and secretory component of immunoglobulins in small intestine of rats treated with *Saccharomyces boulardii*. *Digest Dis Sci* 1990, 35(2):251-256

Buts JP, Cortier G, Delmee M. *Saccharomyces boulardii* for *Clostridium difficile*-associated enteropathies in infants. *J Pediatr Gastroenterol Nutr* 1993, 16:419-425

Buts JP. Twenty-five years of research on *Saccharomyces boulardii* trophic effects: Updates and perspectives. *Digest Dis Sci* 2009, 54(1):15-18

Byron JK, Clemons KV, Mc Cusker JH, Davis RW, Stevens DA. Pathogenicity of *Saccharomyces cerevisiae* in complement factor five-deficient mice. *Infect Immun* 1995, 63:478-485

Can M, Beşirbellioglu BA, Avci IY, Beker CM, Pahsa A. Prophylactic *Saccharomyces boulardii* in the prevention of antibiotic-associated diarrhea: a prospective study. *Med Sci Monit* 2006, 12(4):PI19-22

Canani RB, Cirillo P, Terrin G, Cesarano L, Spagnuolo MI, De Vincenzo A et al. Probiotics for treatment of acute diarrhoea in children: randomized clinical trial of five different preparations. *BMJ* 2007, 335:340-342

Casem RAO. *Saccharomyces boulardii* in the prevention of antibiotic associated diarrhea in children: a randomized controlled trial. *Pediatr Infect Dis Soc Philipp J* 2013, 13(2):70-76

Cassone M, Serra P, Mondello F, Girolamo A, Scafetti S, Pistella E et al. Outbreak of *Saccharomyces cerevisiae* subtype *boulardii* fungemia in patients neighboring those treated with a probiotic preparation of the organism. *J Clin Microbiol* 2003, 41(11):5340-5343

Cetina-Sauri G, Sierra Basto G. Evaluación terapéutica de *S. boulardii* en niños con diarrea aguda. *Comp Inv Lat Am Mex* 1989, 9:108-112 [in Spanish]

Cetina-Sauri G, Sierra Basto G. Antidiarrhöische Therapie bei Kindern-Therapeutische Prüfung von *Saccharomyces boulardii* an Kindern mit acuter Diarröh. *Der Kinderarzt* 1991, 22(12):2059-2061 [in German]

Cetina-Sauri G, Sierra Basto G. Evaluation of *Saccharomyces* for the treatment of acute diarrhea in pediatric patients. *Ann Pediatr* 1994, 41(6):397-400

CFR – Code of Federal regulations Title 21 (FDA), 21CFR172.896. Available at:  
<https://www.accessdata.fda.gov/scripts/cdrh/cfdocs/cfCFR/CFRSearch.cfm?fr=172.896>

Chapoy P. Traitement des diarrhées aiguës infantiles. *Ann Pédiatr* 1985, 32:561-563 [in French]

Chapoy P. Behandlung akuter Diarrhöe bei Kleinkindern. *Therapiewoche* 1986, 40:3-6 [in German]

Chen J, Wan C-M, Gong S-T, Fang F, Sun M, Qian Y et al. Chinese clinical practice guidelines for acute infectious diarrhea in children. *World J Pediatr* 2018, 14:429-436

CHMP (Committee for Medicinal Products for Human use). Guideline on the evaluation of medicinal products for the treatment of irritable bowel syndrome. CPMP/EWP/785/97 Rev. 1 from 25.09.2014

Choi CW, Jo SY, Park HJ, Chang SK, Byeon JS, Myung SJ. A randomized, double-blind, placebo-controlled, multicenter trial of *Saccharomyces boulardii* in irritable bowel syndrome. Effect of quality of life. *J Clin Gastroenterol* 2011, 45:679-683

Chosewood LC, Wilson DE. (editors) Biosafety in Microbiological Laboratories. U.S. Department of Health and Human Services HHS Publication No. (CDC) 21-1112, revised December 2009. Available at: <http://www.cdc.gov/biosafety/publications/bmbl5/index.htm>

Chouraqui JP, Dietsch J, Musial C, Blehaut H. Controlled treatment trial with *Saccharomyces boulardii* (SB) in the irritable-bowel-syndrome of infancy (IBSI) [abstract]. *Gastroenterology* 1995, 4(108):584

Chu Y, Zhu H, Zhou Y, LV L, Huo J. Intervention study on *Saccharomyces boulardii* with proton pump inhibitor (PPI)-based triple therapy for *Helicobacter pylori* related peptic ulcer. *African J Pharm Pharmacol* 2012, 6:2900-2904

Cindoruk M, Erkan G, Karkan T, Dursun A, Unal S. Efficacy and safety of *Saccharomyces boulardii* in the 14-day triple anti-*Helicobacter pylori* therapy: a prospective randomized placebo-controlled double-blind study. *Helicobacter* 2007, 12:309-316

Clemons KV, McCusker JH, Davis RQ, Stevens DA. Comparative pathogenesis of clinical and nonclinical isolates of *Saccharomyces cerevisiae*. *J Infect Dis* 1994, 169:859-867

CMDh: *Saccharomyces boulardii*: CMDh scientific conclusions and grounds for variation, amendments to the product information and timetable for implementation – PSUSA/00009284/201702, last updated: 01 December 2017

Cohen SH, Gerding DN, Johnson S, Kelly CP, Loo VG, McDonald LC et al. Clinical practice guidelines for *Clostridium difficile* infection in adults: 2010 update by the society for healthcare epidemiology of America (SHEA) and the infectious diseases society of America (IDSA). *Infect Control Hosp Epidemiol* 2010, 31:431-455

Corrêa NB, Penna FJ, Lima FM, Nicoli JR, Filho LA. Treatment of acute diarrhea with *Saccharomyces boulardii* in infants. *J Pediatr Gastroenterol Nutr* 2011, 53(5):497-501

Costalos C, Skouteri V, Gounaris A, Sevastiadou S, Triandafilidou A, Ekonomidou C et al. Enteral feeding of premature infants with *Saccharomyces boulardii*. *Early Human Development* 2003, 74:89-96

Cottrell J, Koenig K, Perfekt R, Hofmann R. Comparison of two forms of loperamide-simeticone and a probiotic yeast (*Saccharomyces boulardii*) in the treatment of acute diarrhoea in adults: a randomized non-inferiority clinical trial. *Drugs R D* 2015, 15:363-373, doi 10.1007/s40268-015-0111-y

Cremonini F, DiCaro S, Covino M, Armuzzi A, Gabrielli M, Santarelli L et al. Effect of different probiotic preparations on anti-*Helicobacter pylori* therapy-related side effect: a parallel group, triple blind, placebo-controlled study. *Am J Gastroenterol* 2002a, 97(11):2744-2749

Cremonini F, Di Caro S, Nista EC, Bartolozzi F, Capelli G, Gasbarrini G et al. Meta-analysis: the effect of probiotic administration on antibiotic-associated diarrhea. *Aliment Pharmacol Ther* 2002b, 16:1461-1467

Czerucka D, Rampal P. Diversity of *Saccharomyces boulardii* CNCM I-745 mechanisms of action against intestinal infections. *World J Gastroenterol* 2019, 25(18):2188-2203, doi 10.3748/wjg.v25.i18.2188

Dalgic N, Sancar M, Bayraktar B, Pullu M, Hasim O. Probiotic, zinc and lactose-free formula in children with rotavirus diarrhea: Are they effective? *Pediatr Int* 2011, 53:677-682

Das S, Gupta PK, Das RR. Efficacy and safety of *Saccharomyces boulardii* in acute rotavirus diarrhea: double blind randomized controlled trial from a developing country. *Journal of Tropical Pediatrics* 2016, 62:464-470, doi 10.1093/tropej/fmw032

De Boer WA, Thys JC, Borody TJ, Graham DY, O'Morain C, Tytgat GNJ. Proposal for use of a standard side effect scoring system in studies exploring *Helicobacter pylori* treatment regimens. *Eur J Gastroenterol Hepatol* 1996, 8:641-643

Dendukuri N, Costa V, McGregor M, Brophy M. Probiotic therapy for the prevention and treatment of *Clostridium difficile*-associated diarrhea: a systematic review. *CMAJ* 2005, 173(2):167-170

Didari T, Solki S, Mozaffari S, Nikfar S, Abdollah M. A systematic review of the safety of probiotics. *Expert Opin. Drug Saf.* 2014, 13(2):227-239

Dinleyici EC, Eren M, Yargic ZA, Dogan N, Vandenplas Y. Clinical efficacy of *Saccharomyces boulardii* and metronidazole compared to metronidazole alone in children with acute bloody diarrhea caused by amebiasis: a prospective, randomized, open label study. *Am J Trop Med Hyg* 2009, 80(6):953-955

Dinleyici EC, Eren M, Dogan N, Reyhanioglu S, Yargic ZA, Vandenplas Y. Clinical efficacy of *Saccharomyces boulardii* or metronidazole in symptomatic children with *Blastocystis hominis* infection. *Parasitol Res* 2011, 108:541-545

Dinleyici EC, Eren M, Ozen M, Yargic ZA, Vandenplas Y. Effectiveness and safety of *Saccharomyces boulardii* for acute infectious diarrhea. *Expert Opin Biol Ther* 2012, 12(4):395-410

Duman DG, Bor S, Özütemiz Ö, Sahin T, Oguz D, Isitan F et al. Efficacy and safety of *Saccharomyces boulardii* in prevention of antibiotic-associated diarrhea due to *Helicobacter pylori* eradication. *Eur J Gastroenterol Hepatol* 2005, 17:1357-1361

D'Souza AL, Rajkumar C, Cooke J, Bulpitt CJ. Probiotics in prevention of antibiotic-associated diarrhoea: meta-analysis. *BMJ* 2002, 324:1361-1364

DuPont H. Therapy for and prevention of traveler's diarrhea. *Clin Infect Dis* 2007, 45:S78-84

Edwards-Ingram L, Gitsham P, Burton N, Warhurst G, Clarke I, Hoyle D et al. Genotypic and physiological characterization of *Saccharomyces boulardii* the probiotic strain of *Saccharomyces cerevisiae*. *App Environ Microbiol* 2007, 73(8):2458-2467

Edwards-Ingram LC, Gent ME, Hoyle DC, Hayes A, Stateva LI, Oliver SG. Comparative genomic hybridization provides new insights into the molecular taxonomy of the *Saccharomyces sensu stricto* complex. *Genome Res* 2004, 14(6):1043-1051

EFSA: Scientific Opinion on the substantiation of health claims related to non-characterised bacteria and yeasts pursuant to Article 13(1) of Regulation (EC) No 1924/2006. *EFSA Journal* 2010, 8(2):1470

EFSA: Scientific opinion on the substantiation of health claims related to *Saccharomyces cerevisiae* var. *boulardii* CNCM I-1079 and defense against pathogenic gastro-intestinal microorganisms (ID 913, further assessment) pursuant to article 13(1) of regulation (EC) no 1924/2006. *EFSA Journal* 2012a, 10(6):2717

EFSA: Scientific opinion on the substantiation of a health claim related to *Saccharomyces cerevisiae* var. *boulardii* CNCM I-3799 and reducing gastrointestinal discomfort pursuant to article 13(5) of regulation (EC) no 1924/2006. *EFSA Journal* 2012b, 10(7):2801

Ehrhardt S, Guo N, Hinz R, Schoppen S, May J, Reiser M et al. *Saccharomyces boulardii* to prevent antibiotic-associated diarrhea: a randomized, double-masked, placebo-controlled trial. *Open Forum Infectious diseases* 2016, 3(1), doi 10.1093/ofid/ofw011

Elmer GW, Moyer KA, Vega R, Surawicz CM, Collier AC, Hooton TM et al. Evaluation of *Saccharomyces boulardii* for patients with HIV-related chronic diarrhea and in healthy volunteers receiving antifungals. *Microecol Ther* 1995, 25:23-31

Elmer GW, Martin SW, Horner KL, McFarland LV, Levy RH. Survival of *Saccharomyces boulardii* in the rat gastrointestinal tract and effects of dietary fiber. *Microb Ecology health Dis* 1999a, 37:315-317

Elmer GW, McFarland LV, Surawicz M, Danko L, Greenberg RN. Behaviour of *Saccharomyces boulardii* in recurrent *Clostridium difficile* disease patients. *Aliment Pharmacol Ther* 1999b, 13:1663-1668

Enache-Angoulvant A, Hennequin C. Invasive *Saccharomyces* infection: a comprehensive review. *Clin Infect Dis* 2005, 41:1559-1568

Erdeve O, Tiras U, Dallar Y. The probiotic effect of *Saccharomyces boulardii* in a pediatric age group. *J Trop Pediatr* 2004, 50:234-236

Erdeve O, Tiras U, Dallar Y, Savas S. *Saccharomyces boulardii* and antibiotic-associated diarrhoea in children. *Aliment Pharmacol Ther* 2005, 21:583-590

Erdoğan Ö, Tanyeri B, Torun E, Gönüllü E, Arslan H, Erenberk U, et al. The comparison of the efficacy of two different probiotics in rotavirus gastroenteritis in children. *J Trop Med* 2012, ID:787240

Eren M, Dinleyici EC, Vandenplas Y. Clinical efficacy comparison of *Saccharomyces boulardii* and yogurt fluid in acute non-bloody diarrhea in children: a randomized, controlled, open label study. *Am J Trop Med Hyg* 2010, 82(3):488-491

Eriksson OE, Winka K. Supraordinal taxa of Ascomycota. *Myconet* 1997, 1:1-16

Feizizadeh S, Salehi-Abargouei A, Akbari V. Efficacy and safety of *Saccharomyces boulardii* for acute diarrhea. *Pediatrics* 2014, 134(1):e176-e191

Feng J, Wang F, Qiu X, Mc Farland LV, Chen P, Zhou R et al. Efficacy and safety of probiotic-supplemented triple therapy for eradication of *Helicobacter pylori* in children: a systematic review and network analysis. *Eur J Clin Pharmacol* 2017, 73:1109-1208, doi 10.1007/s00228-017-2291-6

Fischbach W, Malfertheiner P, Hoffmann JC, Bolten W, Bornschein J, Götze O et al. S3-Leitlinie „*Helicobacter pylori* und gastroduodenale Ulkuskrankheit“. *Z Gastroenterol* 2009, 47:68-102 [in German]

Flatley EA, Wilde AM, Nailor MD. *Saccharomyces boulardii* for the prevention of hospital *Clostridium difficile* infection. *Gastrointestinal Liver Dis* 2015, 24(1):21-24, doi 10.15403/jgld.2014.1121.fly

Florez I, Veroniki A, Khalifah R, Yépes-Núñez J, Sierra J, Vernooij R et al. Comparative effectiveness and safety of interventions for acute diarrhea and gastroenteritis in children: a systematic review and network meta-analysis. *PLoS ONE* 2018, 13(12):e0207701

Gao C, Xie R, Ma T, Wu S. Therapeutic effect of *Saccharomyces boulardii* combined with standard triple therapy for *Helicobacter pylori* eradication. *Chin J Gastroenterol* 2012, 17:555-557 [in Chinese, only abstract in English]

Gaon D, Garcia H, Winter L, Rodriguez N, Quintas R, Gonzales SN et al. Effect of *Lactobacillus* strains and *Saccharomyces boulardii* on persistent diarrhea in children. *Medicina (Buenos Aires)* 2003, 63:293-298

Garcia Vilela E, De Lourdes De Abreu Ferrari M, Oswaldo Da Gama Torres H, Guerra Pinto A, Carneiro Aguirre C, Paiva Martins F et al. Influence of *Saccharomyces boulardii* on the intestinal permeability of patients with Crohn's disease in remission. *Scan J Gastroenterol* 2008, 43:842-848

Girard-Pipau F, Pompei A, Schneider S. Intestinal microflora, short chain and cellular fatty acids, influence of a probiotic *Saccharomyces*. *Microb Ecology Health Dis* 2002, 14:220-227

Gorbach SL. Probiotics and gastrointestinal health. *Am J Gastroenterol* 2000, 95:S2-S4

Graff S, Chaumeil J-C, Boy P, Lai-Kuen R, Charrueau Cl. Influence of pH conditions on the viability of *Saccharomyces boulardii* yeast. *J Gen Appl Microbiol* 2008, 54:221-227

Grandy G, Medina M, Soria R, Terán CG, Araya M. Probiotics in the treatment of acute rotavirus diarrhea. A randomized, double-blind, controlled trial using two different probiotic preparations in Bolivian children. *BMC Infect Dis* 2010, 10:253

Guarino A, Ashkenazi S, Gendrel D, Lo Vecchio A, Shamir R, Szajewska H. European Society for Pediatric Gastroenterology, Hepatology, and Nutrition / European Society for pediatric Infectious

Diseases Evidence-based guidelines for the management of acute gastroenteritis in children in Europe: update 2014. *JPGN* 2014, 59(1):132-152

Guarino A, Guandalini S, Lo Vecchio A. Probiotics for prevention and treatment of diarrhea. *J Clin Gastroenterol* 2015, 49:S37-S45

Guo Q, Goldenberg JZ, Humphrey C, El Dib R, Johnston BC. Probiotics for the prevention of pediatric antibiotic-associated diarrhea. *Cochrane Database of Systematic Reviews* 2019, 4(4):CD00482, doi 10.1002/14651858.CD004827.pub5

Hafeez A, Tariq P, Ali S, Kundi ZU, Khan A, Hussan M. The efficacy of *Saccharomyces boulardii* in the treatment of acute watery diarrhea in children: a multicentre randomized controlled trial. *J Coll Physicians Surg Pak* 2002, 12:432-434 [cited in Szajewsak et al. 2007, publication not available]

Hempel S, Newberry SJ, Maher AR, Wang Z, Miles JN, Shanman R et al. Probiotics for the prevention and treatment of antibiotic-associated diarrhea. A systematic review and meta-analysis. *JAMA* 2012, 307(18):1959-1969

Hennequin C, Kaufmann-Lacroix C, Jobert A, Viard JP, Ricour C, Jaquemin JL et al. Possible role of catheters in *Saccharomyces boulardii* fungemia. *Eur J Microbiol Infect Dis* 2000, 19:16-20

Hill DR, Ericsson CD, Pearson RD, Keystone JS, Freedman DO, Kozarsky PE et al. The practice of travel medicine: guidelines by the Infectious Diseases Society of America. *Clin Infect Dis* 2006, 43:1499-1539

Höchter W, Chase D, Hagenhoff G. *Saccharomyces boulardii* bei akuter Erwachsenendiarrhoe. *MMW* 1990, 132:188-192 [in German]

Htwe K, Yee KS, Tin M, Vandenplas Y. Effect of *Saccharomyces boulardii* in the treatment of acute watery diarrhea in Myanmar children: a randomized controlled study. *Am J Trop Med Hyg* 2008, 78(2):214-216

Hurduc V, Plesca D, Dragomir D, Sajin M, Vandenplas Y. A randomized, open trial evaluating the effect of *Saccharomyces boulardii* on the eradication rate of *Helicobacter pylori* infection in children. *Acta Paediatr* 2009, 98(1):127-131

Im E, Pothoulakis C. Recent advances in *Saccharomyces boulardii* research. *Gastroenterol Clin Biol* 2010, 34 (Suppl 1):62-70

Izquierdo-Pulido M, Font-Fábregas J, Vidal-Carou C. Influence of *Saccharomyces cerevisiae* var. *uvarum* on histamine and tyramine formation during beer fermentation. *Food Chem* 1995, 54:51-54

Jahn HU, Ullrich R, Schneider T, Liehr RM, Schieferdecker HL, Holst H et al. Immunological and trophical effects of *Saccharomyces boulardii* on the small intestine in healthy human volunteers. *Digestion* 1996, 57:95-104

Ji D, Zou S, Yuan L. *Saccharomyces boulardii* in treatment of acute children diarrhea: a randomized placebo-controlled trial. *Journal of Pediatric Pharmacy* 2009, 15(1):13-15 [in Chinese, only abstract in English]

Jindal M, Goyal Y, Iata S, Sharma RK. Preventive role of probiotics in antibiotic associated diarrhoea in children. *Indian Journal of Public Health Research & Development* 2017, 8(3):66-69

Johnston BC, Supina AL, Ospina M, Vohra S. Probiotics for the prevention of pediatric antibiotic-associated diarrhea (review). *Cochrane Database of Systematic Reviews* 2007, 2:CD004827

Johnston BC, Goldenberg JZ, Vandvik PO, Sun X, Guyatt GH. Probiotics for the prevention of pediatric antibiotic-associated diarrhea. *Cochrane Database of Systematic Reviews* 2011, 11:CD004827, doi 10.1002/14651858.CD004827.pub3

Kabbani TA, Pallav K, Dowd SE, Villafuerte-Galvez J, Vanga RR, Castillo NE et al. Prospective randomized controlled study on the effects of *Saccharomyces boulardii* CNCM I-745 and amoxicillin-clavulanate or the combination on the gut microbiota of healthy volunteers. *Gut Microbes* 2017, 8(1):17-32

Kabir MA, Ishaque SM, Ali MS, Mahmuduzzaman M, Hasan M. Role of *Saccharomyces boulardii* in diarrhea predominant irritable bowel syndrome. *Mymensingh Med J* 2011, 20(3):397-401

Karen M, Yuksel O, Akyürek N, Ofluoğlu E, Çağlar K, Sahin TT et al. Probiotic agent *Saccharomyces boulardii* reduces the incidence of lung injury in acute necrotizing pancreatitis induced rats. *J Surg Res* 2010, 160(1):139-144

Kelesidis T, Pothoulakis C. Efficacy and safety of the probiotic *Saccharomyces boulardii* for the prevention and therapy of gastrointestinal disorders. *Ther Adv Gastroenterol* 2012, 5(2):111-125

Khan A, Javed T, Chishti AL. Clinical efficacy of use of probiotic "Saccharomyces boulardii" in children with acute watery diarrhea. *Pakistan Paed J* 2012, 36(3):122-127

Kim JH, Lee DH, Jeong SC, Chung KS, Lee JS. Characterizations of antihypertensive angiotensin I-converting enzymes inhibitor from *Saccharomyces cerevisiae*. *J Microbiol Biotechnol* 2004, 14(6):1318-1323

Klein SM, Elmer GW, McFarland LV, Surawicz CM, Levy RH. Recovery and elimination of the biotherapeutic agent, *Saccharomyces boulardii*, in healthy human volunteers. *Pharm Res* 1993, 10(11):1615-1619

Kollaritsch HH, Kremsner P, Wiedermann G, Scheiner O. Prevention of traveller's diarrhea: comparison of different non-antibiotic preparations. *Travel Med Internat* 1989, 7:9-18

Kollaritsch H, Holst H, Grobara P, Wiedermann G. Prophylaxe der Reisediarrhöe mit *Saccharomyces boulardii*. Ergebnisse einer plazebokontrollierten Doppelblindstudie. *Fortschr Med* 1993, 11:152-156 [in German]

Kommission E. Aufbereitungsmonographie Faex medicinalis (Medizinische Hefe). *BAnz*, Nr. 85, 05 May 1988 [in German]

Kommission E. Aufbereitungsmonographie Trockenhefe aus *Saccharomyces cerevisiae* HANSEN CBS 5926 (Synonym: *Saccharomyces cerevisiae*, *BAnz*, Nr. 71, 15 April 1994 [in German]

Korpela R, Niittynen L. Probiotics and irritable bowel syndrome. *Microb Ecol Health Dis* 2012, 23:18573

Kotowska M, Albrecht P, Szajewska H. *Saccharomyces boulardii* in the prevention of antibiotic-associated diarrhoea in children: a randomized double-blind placebo-controlled trial. *Aliment Pharmacol Ther* 2005, 21:583-590

Kujath P, Sipp H. Neuartige Therapiemöglichkeit der Akne vulgaris. *Wehrmed Mschr* 1978, 12:374-376 [in German]

Kurtzman CP, Fell JW. Yeast Systematics and Phylogeny – Implications of Molecular Identification Methods for Studies in Ecology. *The Yeast Handbook Carlos Augusto Rosa, Gabor Peter, Springer-Verlag*, Berlin Heidelberg 2006

Kurugöl Z, Koturoglu G. Effects of *Saccharomyces boulardii* in children with acute diarrhoea. *Acta Paediatr* 2005, 94:44-47

Kyriakos N, Papamichael K, Roussos A, Theodoropoulos I, Karakoidas C, Smyrnidis A et al. A lyophilized form of *Saccharomyces boulardii* enhances the *Helicobacter pylori* eradication rates of omeprazole-triple therapy in patients with peptic ulcer disease or functional dyspepsia. *Hospital Chronicles* 2013, 8(3):127-133

Lacarrière C, Rieckhof B. Behandlung akuter Diarrhöe in der Praxis. *Therapiewoche* 1986, 36:2551-2556 [in German]

Laux G, Ulrich S. Tranylcypromin. *PPT* 2006, 13 (4):130-41 [in German]

Lee J, Choi SP, Park JH, Shin JW, Joo YH. The effect of *Saccharomyces boulardii* as an adjuvant to the 14-day triple therapy for eradication of *Helicobacter pylori* [abstract]. *J Gastroenterol Hepatol* 2011, 26(5):257

Le Luyer B, Makhoul G, Duhamel JF. Étude multicentrique, contrôlée en double insu d'une formule adaptée enrichie en *Saccharomyces boulardii* dans le traitement des diarrhées aiguës du nourrisson. *Arch Pédiatr* 2010, 17:459-465 [in French]

Lentze MJ. Physiologie des Verdauungstraktes im Kindesalter. In: von Schweinitz D, Ure B (eds.) Kinderchirurgie. Springer-Verlag, Berlin Heidelberg 2013

Lewis SJ, Potts LF, Barry RE. The lack of therapeutic effect of *Saccharomyces boulardii* in the prevention of antibiotic-related diarrhea in elderly patients. *J Infect* 1998, 36:171-174

Lherm T, Monet C, Nougière B, Soulier M, Larbi D, Le Gall C et al. Seven cases of fungemia with *Saccharomyces boulardii* in critically ill patients. *Intensive Care Med* 2002, 28(6):797-801

Li BZ, Threapleton DE, Wang J-Y, Xu JM, Yuan JQ, Zhang C et al. Comparative effectiveness and tolerance of treatments for *Helicobacter pylori*: systematic review and network meta-analysis. *BMJ* 2015, 351:h4052

Lukaszewicz M (2012) *Saccharomyces cerevisiae* var. *boulardii* – Probiotic Yeast. In: Rigobelo EC (ed.), Probiotics. Chapter 16. InTech 2012

Machado Caetano JA, Paramés MT, Babo MJ, Santos A, Bandeira Ferreira, Freitas AA et al. Immunopharmacological effects of *Saccharomyces boulardii* in healthy human volunteers. *Int J Immunopharmac* 1986, 8(3):245-259

Maejima K, Shimoda K, Morita C, Fujiwara T, Kitamura T. Colonization and pathogenicity of *Saccharomyces cerevisiae*, MC 16, in mice and cynomolgus monkeys after oral and intravenous administration. *Jpn J Med Sci Biol* 1980, 33(5):271-276

Malgoire JY, Bertout S, Renaud F, Bastide JM, Mallié M. Typing of *Saccharomyces cerevisiae* clinical strains by using microsatellite sequence polymorphism. *J Clin Microbiol* 2005, 43(3):1133-1137

Mandrella C. Erfolgreicher Behandlungsversuch der Akne mit einem neuartigen Hefepräparat. *Die Heilkunst* 1973, 2:48-50 [in German]

Mansour-Ghanaei F, Dehbashi N, Yazdanparast K, Shafaghi A. Efficacy of *Saccharomyces boulardii* in acute amoebiasis. *World J Gastroenterol* 2003, 9:1832-1833

Marshall BJ, Warren JR. Unidentified curved bacilli in the stomach of patients with gastritis and peptic ulceration. *Lancet* 1984, 16:1311-1314

Martin SW, Heatherington AC, Elmer GW. Pharmacokinetics of Biotherapeutic Agents. In: Elmer GW, McFarland LV, Surawicz CM (eds.) *Biotherapeutic Agents and Infectious Diseases*. Chapter 3. Humana Press Inc Totowa NJ 1999:47-84

Maupas JL, Champemont P, Delforge M. Treatment of irritable bowel syndrome. Double-blind trial of *Saccharomyces boulardii*. *Med Chirur Dig* 1983, 12(1):77-79

Mayr P, Hotz J, Brandau J, Körner J. Erfolgreiche Durchfallbehandlung mit *Saccharomyces boulardii*. *Der Kassenarzt* 1996, 43:52-60 [in German]

McCullough MJ, Clemons KV, Cusker JH, Stevens DA. Species identification and virulence attributes of *Saccharomyces boulardii* (nom. inval.). *J Clin Microbiol* 1998, 36:2613-2617

McFarland LV, Surawicz CM, Greenberg RN, Fekety R, Elmer GW, Moyer KA et al. A randomized placebo-controlled trial of *Saccharomyces boulardii* in combination with standard antibiotics for *Clostridium difficile* disease. *JAMA* 1994, 271:1913-1918

McFarland LV, Surawicz CM, Greenberg RN, Elmer GW, Moyer KA, Melcher SA et al. Prevention of β-lactam associated diarrhea by *Saccharomyces boulardii* compared with placebo. *Am J Gastroenterol* 1995, 90(3):439-448

McFarland LV. *Saccharomyces boulardii* is not *Saccharomyces cerevisiae*. *Clin Infect Dis* 1996, 22:200-201

McFarland LV. Meta-analysis of probiotics for the prevention of antibiotic associated diarrhea and the treatment of *Clostridium difficile* disease. *Am J Gastroenterol* 2006, 101:812-822

McFarland LV. Meta-analysis of probiotics for the prevention of traveler's diarrhea. *Travel Med Infect Dis* 2007, 5:97-105

McFarland LV, Dublin S. Meta-analysis of probiotics for the treatment of irritable bowel syndrome. *World J Gastroenterol* 2008, 14(17):2650-2661

McFarland LV. Systematic review and meta-analysis of *Saccharomyces boulardii* in adult patients. *World J Gastroenterol* 2010, 16(18):2202-2222

Micklefield G. *Saccharomyces boulardii* bei Antibiotika-assozierter Diarrhoe. *MMW Fortschr Med Originalien* 2014, 156(Suppl. 1):18-22 [in German]

Mitterdorfer G, Mayer HK, Kneifel W, Viernstein H. Clustering of *Saccharomyces boulardii* strains within the species *S. cerevisiae* using molecular typing techniques. *J Appl Microbiol* 2002a, 93(4):521-530

Mitterdorfer G, Mayer HK, Kneifel W, Viernstein H. Protein fingerprinting of *Saccharomyces* isolates with therapeutic relevance using one-and two-dimensional electrophoresis. *Proteomics* 2002b, 2(11):1532-1538

Monteiro E, Fernandes JP, Vieira MR, Correia JP, Caetano JM, Ribeiro T et al. Ensaio clinico duplamente cego da ultra-levure na profilaxia das perturbações gastro-intestinais e muco-cutâneas da antibioticaterá. *Acta Med Port* 1981, 3:143-145 [in Portuguese]

Moré M, Swidsinski A. *Saccharomyces boulardii* CNCM I-745 supports regeneration of the intestinal microbiota after diarrheic dysbiosis – a review. *Clin Exp Gastroenterol* 2015, 8:237-255

Moré M, Vandenplas Y. *Saccharomyces boulardii* CNCM I-745 Improves Intestinal Enzyme Function: A Trophic Effects Review. *Clin Med Insights Gastroenterol* 2018, 11:1-14

Moslehi-Jenabian S, Pedersen LL, Jespersen L. Beneficial effects of probiotic and food borne yeasts on human health. *Nutrients* 2010, 2:449-473

Ozkan TB, Sahin E, Erdemir G, Budak F. Effect of *Saccharomyces boulardii* in children with acute gastroenteritis and its relationship to the immune response. *J Int Med Res* 2007, 35:201-212

Padayachee M, Visser J, Viljoen E, Musekiwa A, Blaauw R. Efficacy and safety of *Saccharomyces boulardii* in the treatment of acute gastroenteritis in the paediatric population: a systematic review. *South African Journal of Clinical Nutrition* 2019, 32(3):58-69, doi 10.1080/16070658.2018.1449378

Pan J, Hu J, Qi X, Tu Z. A meta analysis of the efficacy of *Saccharomyces boulardii* in children with acute diarrhea. *Afr J Pharm Pharmacol* 2012, 6(21):1508-1514

Pecquet S, Guillaumin D, Tancrede C, Andremont A. Kinetics of *Saccharomyces cerevisiae* elimination from the intestines of human volunteers and effect of this yeast on resistance to microbial colonization in gnotobiotic mice. *Appl Environ Microbiol* 1991, 57(10):3049-3051

Perapoch J, Planes AM, Querol A, López V, Martínez-Bendayán I, Tormo R et al. Fungemia with *Saccharomyces cerevisiae* in two newborns, only one of whom had been treated with ultra-levura. *Eur J Clin Microbiol Infect Dis* 2000, 19(6):468-470

Piarroux R, Millon L, Bardouet K, Vagner O, Koenig H. Are live *Saccharomyces* yeast harmful to patients? *Lancet* 1999, 353:1851-1852

Pineton de Chambrun G, Neut C, Chau A, Cazaubiel M, Pelerin F, Justen P et al. A randomized clinical trial of *Saccharomyces cerevisiae* versus placebo in the irritable bowel syndrome. *Digest Liver Dis* 2015, 47:119-124

Pozzoni P, Riva A, Bellatorre AG, Amigoni M, Redaelli E, Ronchetti A et al. *Saccharomyces boulardii* for the prevention of antibiotic-associated diarrhea in adult hospitalized patients: a single-center, randomized, double-blind, placebo-controlled trial. *Am J Gastroenterol* 2012, 107:922-931

Reuter J, Merfort I, Schempp CM. Botanicals in dermatology. An evidence-based review. *Am J Clin Dermatol* 2010, 11(4):247-267

Riaz M, Alam S, Malik A, Ali SM. Efficacy and safety of *Saccharomyces boulardii* in acute childhood diarrhea: a double blind randomized controlled trial. *Indian J Pediatr* 2012, 79(4):478-482

Riquelme AJ, Calvo MA, Guzmán AM, Depix MS, García P, Pérez C et al. *Saccharomyces cerevisiae* fungemia after *Saccharomyces boulardii* treatment in immunocompromised patients. *J Clin Gastroenterol* 2003, 36(1):41-43

Roy U, Jessani LG, Rudramurthy SM, Gopalakrishnan R, Dutta S, Chakravarty C et al. Seven cases of *Saccharomyces* fungaemia related to use of probiotics. *Mycoses* 2017, 60:375-380, in press, doi 10.1111/myc.12604

Saint-Marc T, Blehaut H, Musial C, Touraine JL. Diarrhées en relation avec le sida. *Sem Hôp Paris* 1995, 71(23-24):735-741 [in French]

Sankoff D. Reconstructing the history of yeast genomes. *PLoS Genet* 2009, 5:e1000483

Savaş-Erdeve S, Gökyay S, Dallar Y. Efficacy and safety of *Saccharomyces boulardii* in amebiasis-associated diarrhea in children. *Turk J Pediatr* 2009, 51(3):220-224

Scevola D, Perversi L, Cavanna C, Candiani C, Uberti F, Castiglioni B et al. Acid tolerance and fecal recovery following oral administration of *Saccharomyces cerevisiae*. *J Chemoth* 2003, 15(2):143-147

Schlotterer M, Bernasconi P, Lebreton F, Wassermann D. Intérêt de *Saccharomyces boulardii* dans la tolérance digestive de la nutrition entérale à débit continu chez le brûlé. *Nutr Clin Metabol* 1987, 1:31-34 [in French]

Schneider SM, Girard-Pipau F, Filippi J, Hébutterne X, Moyse D, Hinojosa GC et al. Effects of *Saccharomyces boulardii* on fecal short chain fatty acids and microflora in patients on long-term total enteral nutrition. *World J Gastroenterol* 2005, 11:6165-6169

Shaikh F, Bhand SA, Kumar P, Nizamani MA. Acute diarrhea; role of *Saccharomyces boulardii* in the treatment. *Professional Med J* 2015, 22(3):281-284

Shan LS, Hou P, Wang ZJ, Liu FR, Chen N, Shu LH et al. Prevention and treatment of diarrhea with *Saccharomyces boulardii* in children with acute lower respiratory tract infections. *Beneficial Microbes* 2013, 4(4):329-334

Shen Y, Liu W, Jiang L, Shao C, Xu C. Effects of *Saccharomyces boulardii* in children with acute diarrhea. *J Clin Pediatr* 2008, 26(6):528-531 [in Chinese, only abstract in English]

Song J, Park DI, Park JH, Kim HJ, Cho YK, Sohn CI et al. The effect of probiotics and mucoprotective agents on PPI-based triple therapy for eradication of *Helicobacter pylori*. *Helicobacter* 2010, 15:206-213

Summary of Product Characteristics (SmPC), various products

Stickl H. Das Immunverhalten des Menschen nach Einnahme von Hefe. *Natura-med* 1987, 3:156-157 [in German]

Stüttgen M. Aknetherapie mit *Saccharomyces boulardii*. *Akt Dermatol* 1991, 17:206-209 [in German]

Suardi E, Crippa F, d'Arminio Montforte A. Probiotics in the prevention of antibiotic-associated diarrhea in adults. *Int J Probiotics Prebiotics* 2013, 8(1):41-44

Sudha MR. Safety assessment studies of probiotic *Saccharomyces boulardii* strain Unique 28 in Sprague-Dawley rats. *Benef Microbes* 2011, 2(3):221-227

Surawicz CM, Elmer GW, Speelman P, McFarland L, Chinn J, van Belle G. Prevention of antibiotic-associated diarrhea by *Saccharomyces boulardii*: a prospective study. *Gastroenterology* 1989a, 96:981-988

Surawicz CM, McFarland LV, Elmer G, Chinn J. Treatment of recurrent *Clostridium difficile* colitis with vancomycin and *Saccharomyces boulardii*. *Am J Gastroenterol* 1989b, 84(19):1285-1287

Surawicz CM, McFarland L, Greenberg RN, Rubin M, Fekety R, Mulligan RN et al. The search for a better treatment for recurrent *Clostridium difficile* disease: use of high-dose vancomycin combined with *Saccharomyces boulardii*. *Clin Infect Dis* 2000, 31:1012-1017

Swidsinski A, Loening-Baucke V, Verstralen H, Osowska S, Doerffel Y. Biostructure of fecal microbiota in healthy subjects and patients with chronic idiopathic diarrhea. *Gastroenterology* 2008, 135(2):568-579

Swidsinski A, Loening-Baucke V, Schulz S, Manowsky J, Verstraeten H, Swidsinski S. Functional anatomy of the colonic bioreactor: Impact of antibiotics and *Saccharomyces boulardii* on bacterial composition in human fecal cylinders. *Systematic and Applied Microbiology* 2016, 39:67-75, doi 10.1016/j.syamp.2015.11.002

Szajewska H, Mrukowicz J. Meta-analysis: non-pathogenic yeast *Saccharomyces boulardii* in the prevention of anti-biotic-associated diarrhea. *Aliment Pharmacol Ther* 2005, 22:365-372

Szajewska H, Ruszcynski M, Radzikowski A. Probiotics in the prevention of antibiotic-associated diarrhea in children: a meta-analysis of randomized controlled trials. *J Pediatr* 2006, 149:367-372

Szajewska H, Skórka A, Dylag M. Meta-analysis: *Saccharomyces boulardii* for treating acute diarrhoea in children. *Aliment Pharmacol Ther* 2007, 25:257-264

Szajewska H, Skórka A. *Saccharomyces boulardii* for treating acute gastroenteritis in children: updated meta-analysis of randomized controlled trials. *Aliment Pharmacol Ther* 2009, 30:955-963

Szajewska H, Horvath A, Piwowarczyk A. Meta-analysis: the effects of *Saccharomyces boulardii* supplementation on *Helicobacter pylori* eradication rates and side-effects during treatment. *Aliment Pharmacol Ther* 2010, 32:1069-1079

Szajewska H, Guarino A, Hojsak I, Indrio F, Kolacek S, Shamir R et al. Use of probiotics for management of acute gastroenteritis: a position paper by the ESPGHAN Working Group for Probiotics and Prebiotics. *JPGN* 2014, 58(4):531-539

Szajewska H, Horvath A, Kolodziej M. Systematic review with meta-analysis: *Saccharomyces boulardii* supplementation and eradication of *Helicobacter pylori* infection. *Aliment Pharmacol Ther* 2015a, 41:1237-1245

Szajewska H, Kolodziej M. Systematic review with meta-analysis: *Saccharomyces boulardii* in the prevention of antibiotic-associated diarrhea. *Aliment Pharmacol Ther* 2015b, 42:793-801

Tempé JD, Steidel AL, Blehaut, Hasselmann M, Lutun P, Maurier F. Prévention par *Saccharomyces boulardii* des diarrhées de l'alimentation entérale à débit continu. *Sem Hop Paris* 1983, 59(18):1409-1412 [in French]

Terciolo C, Dapoigny M, Andre F. Beneficial effects of *Saccharomyces boulardii* CNCM I-745 on clinical disorders associated with intestinal barrier disruption. *Clin Exper Gastroenterol* 2019, 12:67-82

Tung JM, Dolovich LR, Lee CH. Prevention of *Clostridium difficile* infection with *Saccharomyces boulardii*: a systematic review. *Can J Gastroenterol* 2009, 23(12):817-821

Turck D, Bernet J-P, Marx J, Kempf H, Giard P, Walbaum O et al. Incidence and risk factors of oral antibiotic-associated diarrhea in an outpatient pediatric population. *J Pediatr Gastroenterol Nutr* 2003, 37:22-26

Urgancı N, Polat T, Uysalol M, Çetinkaya F. Evaluation on the efficacy of *Saccharomyces boulardii* in children with acute diarrhoea. *Arch Gastroenterohepatol* 2001, 20(3-4):81-83

Vandenplas Y, Badriul H, Thapa B, Elizabeth K, Bhave S. A multi-centre DBRPC-trial in developing countries with *S. boulardii* in acute gastroenteritis. *J Pediatr Gastroenterol Nutr* 2007, 44(6):e86

Vandenplas Y, Brunser O, Szajewska H. *Saccharomyces boulardii* in childhood. *Eur J Pediatr* 2008, 168(3):253-265

Vaughan-Martini A, Martini A. Three newly delimited species of *Saccharomyces sensu strictu*. *Antonie van Leeuwenhoek* 1987, 53:77-87

Vaughan-Martini A. Reflections on the Classification of yeast for different end-users in biotechnology, ecology, and medicine. *Int Microbiol* 2003, 6:175-182

Venugopalan V, Shriner KA, Wong-Beringer A. Regulatory oversight and safety of probiotic use. *Emerg Infect Dis* 2010, 16:1661-1665

Villarruel G, Rubio DM, Lopez F, Cintioni J, Gurevech R, Romero G et al. *Saccharomyces boulardii* in acute childhood diarrhoea: a randomized, placebo-controlled study. *Acta Paediatr* 2007, 96(4):538-541

Weber G, Adamczyk A, Freytag S. Behandlung der Akne mit einem Hefepräparat. *Fortschr Med* 1989, 10:563-566 [in German]

Xie C, Li J, Wang K Li Q, Chen D. Probiotics for the prevention of antibiotic-associated diarrhea in older patients: a systematic review. *Travel Med Infect Dis* 2015, 13(2):128-134, doi 10.1016/j.tmaid.2015.03.001

Zhao HM, OU-Yang HJ, Duan BP, Xu B, Chen ZY, Tang J et al. Clinical effect of triple therapy combined with *Saccharomyces boulardii* in the treatment of *Helicobacter pylori* infection in children. *Chin J Contemp Pediatr* 2014, 16(3):230-233 [in Chinese, only abstract in English]

Zojaji H, Ghobakhlou M, Rajabalinia H, Ataei E, Sherafat SJ, Moghimi-Dehkordi B. The efficacy and safety of adding the probiotic *Saccharomyces boulardii* to standard triple therapy for eradication of *H. pylori*: a randomized controlled trial. *Gastroenterol Hepatol* 2013, 6(1):S99-S104

#### **References assessed but not cited in the assessment report**

FAO/WHO Expert Consultation Group. Probiotics in Food: Health and nutritional properties and guidelines for evaluation. Nutrition Paper 85, ISSN 0254-4725; Rome 2006; Available at: <https://www.fao.org/3/a0512e/a0512e.pdf>

LPT. Mutagenicity study of *Saccharomyces cerevisiae* HANSEN CBS 5926 in the *Salmonella typhimurium* reverse mutation assay (*in vitro*). 2014. Report No. 30428/1